



**Centurion**  
**UNIVERSITY**

*Shaping Lives...  
Empowering Communities...*

# Chemistry

## SUB. CODE- BSCC2402

*For B.Sc. 4<sup>th</sup> Semester*

### **(Heterocycles)**

### **Module-IV**

***Quinoline and Isoquinoline: structure elucidation***

***By***

***Dr. Rosy Mallik***

***Assistant Professor,***

***CENTURION UNIVERSITY OF TECHNOLOGY AND MANAGEMENT,  
ODISHA***




Centurion  
UNIVERSITY

Shaping Lives...  
Empowering Communities...

# Quinoline: structure elucidation

1. **Molecular formula** is  $C_9H_7N$  deduced from analytical and molecular weight determinations
2. It undergoes electrophilic aromatic substitution reactions like nitration, sulphonation, etc. but does not give Friedel-Craft reaction.
3. It is resistant to addition reactions.
4. On vigorous oxidation with alk.  $KMnO_4$ , it gives pyridine-2,3-dicarboxylic acid (quinolic acid) and oxalic acid.







**Centurion**  
**UNIVERSITY**  
Shaping Lives...  
Empowering Communities...

# Quinoline: structure elucidation

From the reactions, the two possible structures for quinolone are proposed as A or B




**A**



**B**

Quaternary ammonium salts are formed with alkyl halides. This suggests the presence of a tertiary N-atom in the ring.

Hence the pyridine ring should be fused with a benzene ring through a common carbon pair in quinolone.





Centurion  
UNIVERSITY

Shaping Lives...  
Empowering Communities...

# Isoquinoline: structure elucidation

1. **Molecular formula** is  $C_9H_7N$  deduced from analytical and molecular weight determinations
2. It undergoes electrophilic aromatic substitution reactions like nitration, sulphonation, etc. as well as nucleophilic substitution reaction similar to pyridine. This shows the presence of benzene and pyridine ring.
3. It undergoes addition reaction with only one mole of  $CH_3I$  to give quaternary ammonium salt which shows the presence of one tertiary N-atom..




**Centurion**  
**UNIVERSITY**

Shaping Lives...  
Empowering Communities...

# Isoquinoline: structure elucidation

4. On oxidation with alk.  $\text{KMnO}_4$ , it gives mixture of phthalic acid and cinchomeronic acid.



This shows the pyridine ring is not stable towards oxidation, and isoquinoline consists of one benzene ring fused with pyridine ring at 3, 4- position. Hence the structure can be proposed as



isoquinoline




Centurion  
UNIVERSITY

Shaping Lives...  
Empowering Communities...

# Isoquinoline: structure elucidation

The structure is further confirmed by Bischler-Napieralski synthesis





**Centurion**  
**UNIVERSITY**

*Shaping Lives...*  
*Empowering Communities...*

# References

1. **K.D Sharma and Y. R. Sharma, Kalyani Publishers, Unit IV**
2. **University Chemistry, Vol-IV, Dr. U. N. Dash, Dr. K. K. Ojha, Himalaya Publishing house, Unit V**