



Centurion
UNIVERSITY
*Shaping Lives...
Empowering Communities...*

REDOX TITRATIONS - CONCEPTS AND PRINCIPLE

BY, LIPSA SAMAL

ASST. PROF (PA & QA), SPLS, CUTM



Centurion
UNIVERSITY

*Shaping Lives...
Empowering Communities...*

- **Oxidation-reduction reactions involve a transfer of electrons.**
- **The oxidising agent accepts electrons and the reducing agent gives electrons.**
- **In working out the equation for a redox reaction we can use the half-equation method. this method we use the half-equation for the oxidising agent and the half-equation for the reducing agent then add them together.**



Centurion
UNIVERSITY

*Shaping Lives...
Empowering Communities...*

FORMAL POTENTIALS

- **Standard potentials E^0 are evaluated with full regard to activity effects and with all ions present in simple form: they are really limiting or ideal values and are rarely observed in a potentiometric measurement.**

$E = E' + 0.0591/n \cdot \log [Ox] / [Red]$ at 25 degree celcius.

where E' is the formal potential and corresponds to the value of E at unit concentrations of oxidant and reductant, and the quantities in square brackets refer to molar concentrations



Centurion
UNIVERSITY

Shaping Lives...

Empowering Communities...

Internal oxidation-reduction indicators.

acid-base indicators are employed to mark the sudden change in pH during acid - base titrations. Similarly an oxidation- reduction indicator should mark the sudden change in the oxidation potential in the neighbourhood of the equivalence point in an oxidation-reduction titration. The ideal oxidation-reduction indicator will be one with an oxidation potential intermediate between that of the solution titrated and that of the titrant, and which exhibits a sharp, readily detectable colour change. An oxidation-reduction indicator (redox indicator) is a compound which exhibits different colours in the oxidised and reduced forms: $\text{In}_{\text{ox}} + n e \longrightarrow \text{In}_{\text{Red}}$



Centurion
UNIVERSITY

*Shaping Lives...
Empowering Communities...*

- **The oxidation and reduction should be reversible. At a potential E the ratio of the concentrations of the two forms is given by the Nernst equation:**
- **$E = E' + 0.0591/n \cdot \log [Ox] / [Red]$**
- **where E' is the standard (strictly the formal) potential of the indicator. If the colour intensities of the two forms are comparable a practical estimate of the colour-change interval corresponds to the change in the ratio $[Inox]/[InRed]$ from 10 to /0' this leads to an interval.**



Centurion
UNIVERSITY

Shaping Lives...
Empowering Communities...

- **If the colour intensities of the two forms differ considerably the intermediate colour is attained at potential somewhat removed from E_{\sim} , but the error is unlikely to exceed 0.06 volt. For a sharp colour change at the end point, $E_{t;}$; should differ by about at least 0.15 volt from the standard (formal) potentials of the other systems involved in the reaction.**



Centurion
UNIVERSITY

Shaping Lives...
Empowering Communities...

THANK YOU