

Algae, Protozoa, Yeasts, Viruses and Fungi

General Characteristics



Algae - General Characteristics

- Occur in abundance in oceans, seas, salt and fresh water lakes, ponds, and streams
- Small aquatic forms make up a large part of the water called plankton
- Phytoplankton is made up of plants and grow where there is sufficient light, moisture, and nutrients
- Have a wide range of size and shape, can occur as single cells that are spherical, rod or spindle shaped
- Algal cells are eukaryotic with thin and rigid cell wall



<u>Euglena - General</u> Characteristics

- Motile algae such as *Euglena* have flexible cell membranes called periplasts
- Cells walls are surrounded by a flexible, gelatinous outer matrix secreted through the cell wall
- Algae contain discrete nucleus, starch grains, oil droplets, vacuoles
- Chlorophyll and other pigments are found in membrane bound organelles known as chloroplasts



<u>Algae - General</u> <u>Characteristics</u>

- Within the choloroplast matrix are found flattened membranous vesicles called thylakoids
- Three types of photosynthetic pigments are found in algae: chlorophylls, carotenoids, and biloproteins
 - Chlorophylls a (found in photosynthetic organisms) b, c, d, e (very rare and found only in two genera)
 - Carotenoids lipid soluble (carotenes, xanthophylls)
 - Biloproteins (phycobilins) water soluble, pigment-protein complexes are present



Euglena





<u>Protozoa</u>

- Found in all moist habitats (sea, fresh water, soil)
- Size and shape show considerable variation Leishmania donovani (small), Amoeba proteus (large)
- Single cells without cell walls
- Intracellular structures consists of cytoplasm separated from the surrounding medium by a cell envelope, nucleus/nuclei
- Cytoplasm is differentiated into ectoplasm (gel like) and endoplasm (fluid and voluminous), other structures include mitochondria, endoplasmic reticulum, Golgi complex, food vacuoles, nuclei



<u>Protozoa</u>

- The cell has at least one eukaryotic nucleus with many protozoa having multiple nuclei
- Structural elements of the nucleus include chromosomes, nuclear membrane, karyoplasm (nucleoplasm)
- Cytoplasm is separated from external environment by cell membrane (plasmalemma) with compound coverings of membranes referred to as pellicle
- Many protozoa form resistant cysts at certain times of their life cycle



<u>Protozoa</u>

- Move by three types of specialized organelles: pseudopodia (temporary projection of part of the cytoplasm), flagella (fine filamentous extension of the cell), and cilia (fine short thread like extensions from the cell)
- Multiply by asexual reproduction (simple binary fission), multiple fission, budding (motile offspring)
- Capacity to regenerate lost parts is characteristic of all protozoa



Protozoan Cell Structure





<u>Viruses</u>

□ Viruses are not cells

- Infectious particles consisting of a nucleic acid enclosed in a protein coat or membranous envelope
- □ The capsid is the protein shell/coat enclosing the genome
- Capsids are built by a large number of protein subunits called capsomeres
- Most complex capsids are found in viruses that infect bacteria – bacteriophages or phages
- □ A complete virus particle is called virion



<u>Viruses</u>

- The smallest genomes exhibiting greatest variation
- Some are DNA whereas others are RNA
- Can be double (ds) or single stranded (ss)
- Most dsDNA genomes contain only a single molecule (monopartite)







DNA Viruses

- Typically classified as 'small' or 'large' genomes
- Use nearly all of the DNA for encoding proteins
- Gene clustering is a feature of small genome viruses
- Small genomes use host DNA polymerase for replication
- Large genomes encode a DNA polymerase for replication



RNA Viruses

- **Encode limited number of proteins**
- Some ssRNA viruses use RNA to encode genes and are referred to as positive-strand ssRNA genomes
- Often encoded protein is RNAdependent RNA polymerase
- Monopartite ssRNA viruses encode a single polyprotein which is further broken down into small molecules



Viruses - Infectivity

- □ Are obligate intracellular parasites
- □ Can reproduce only within a host cell
- Lack the enzymes necessary for metabolism and ribosomes for protein synthesis
- Can infect only a limited range of host cells - host range
- Identify host cells by a "lock and key" fit



Reproduction

- Inside the host the viral genome programs the cell to copy nucleic acid and produce proteins
- Host provides nucleotides, ribosomes, tRNAs, amino acids, ATP, and other components
- DNA viruses use the DNA polymerase of the host cell and RNA viruses use virus-encoded polymerases that use RNA as template
- □ The nucleic acid and capsomeres assemble into viral particles and exit the host cell







<u>Fungi – General</u> <u>Characteristics</u>

- Fungi are eukaryotic chaemoorganotrophic organisms that have no chlorophyll
- The thallus consists of filaments that are commonly branched
- Thallus consists of two parts mycelium (complex of several filaments called hyphae) spores
- There are three types of hyphae nonseptate (coenocytic), septate with uninucleate cells and septate with multinucleate cells
- Some fungi are dimorphic



<u>Fungi</u>

- **Reproduction:** Asexual also called somatic or vegetative reproduction
 - Fission of somatic cells giving rise to two similar daughter cells
 - Budding of somatic cells small outgrowth of parental cell developing into a new individual
 - Fragmentation of the hyphal cells, with each fragment becoming a new organism
 - Spore formation

Molds – Asexual spores

enturion

Empowering Communities...

ng Lives..

VERSITY

Sporangiospores – single celled spores formed in sporangia at the end of special hyphae

- Conidiospores small, single celled conidia formed at the tip of hypha
- Oidia single celled spores formed by disjointing of hyphal cells
- Chlamydospores thick walled single celled spores formed from the cells of vegetative hypha
- Blastospores spores formed by budding



<u>Molds</u>

- Sexual reproduction is by fusion of the compatible nuclei of two parent cells
- Sexual spores produced by fusion of two nuclei occur less frequently
 - Ascospores single celled spores (8) produced in ascus
 - Basidiospores single celled spores borne on club-shaped structure called a basidium
 - Zygospores large thick walled spores formed when the tips of two compatible hyphae combine
 - Oospores formed within a special female structure, the oogonium





<u>Yeasts – General</u> <u>Characterisitcs</u>

- Yeasts are usually unicellular, larger than most bacteria
- Vary considerably in size, commonly egg shape, but some are elongated and some spherical
- Have no flagella or other organs of locomotion
- Consists of a protoplast contained in a cell wall (thick)
- Numerous granules and vacuoles are found



<u>Yeasts – General</u> <u>Characterisitcs</u>

- Reproduction is by budding, by fission, or by a combination of these two processes
 - Budding cell wall softens at one point and protoplasm bulges out
 - Fission similar to bacteria but unlike bacteria they develop a cross wall
 - Sexual Reproduction reproduce by intracellular spore formation



YEAST CELL

