



**Centurion**  
**UNIVERSITY**

*Shaping Lives...  
Empowering Communities...*

# **Centurion University of Technology And Management**

Mathematical Physics-3,  
Dr.Pratibha Tripathi,

BSc Physics Sem-4,2019.  
School of Applied Sciences.



**Centurion**  
**UNIVERSITY**

*Shaping Lives...*  
*Empowering Communities...*

# Module Division

Module 1,2,3- Complex Analysis

Module 4,5,6,7 - Integral Transforms



**Centurion**  
**UNIVERSITY**

*Shaping Lives...*  
*Empowering Communities...*

# **Module 1 , 2 , 3**

## **Complex Analysis**



**Centurion**  
**UNIVERSITY**

Shaping Lives...  
Empowering Communities...

## Definitions

**Definition 1.1** Complex numbers are defined as ordered pairs  $(x, y)$

Points on a complex plane. Real axis, imaginary axis, purely imaginary numbers. Real and imaginary parts of complex number. Equality of two complex numbers.

**Definition 1.2** The sum and product of two complex numbers are defined as follows:

$$\begin{aligned}(x_1, y_1) + (x_2, y_2) &= (x_1 + x_2, y_1 + y_2) \\ (x_1, y_1) \cdot (x_2, y_2) &= (x_1x_2 - y_1y_2, x_1y_2 + x_2y_1)\end{aligned}$$

In the rest of the chapter use  $z, z_1, z_2, \dots$  for complex numbers and  $x, y$  for real numbers. introduce  $i$  and  $z = x + iy$  notation.

## Algebraic Properties

### 1. Commutativity

$$z_1 + z_2 = z_2 + z_1, \quad z_1 z_2 = z_2 z_1$$

### 2. Associativity

$$(z_1 + z_2) + z_3 = z_1 + (z_2 + z_3), \quad (z_1 z_2) z_3 = z_1 (z_2 z_3)$$

### 3. Distributive Law

$$z(z_1 + z_2) = z z_1 + z z_2$$

### 4. Additive and Multiplicative Identity

$$z + 0 = z, \quad z \cdot 1 = z$$

### 5. Additive and Multiplicative Inverse

$$\begin{aligned}-z &= (-x, -y) \\ z^{-1} &= \left( \frac{x}{x^2 + y^2}, \frac{-y}{x^2 + y^2} \right), \quad z \neq 0\end{aligned}$$



**Centurion**  
**UNIVERSITY**

Shaping Lives...  
Empowering Communities...

#### 6. Subtraction and Division

$$z_1 - z_2 = z_1 + (-z_2), \quad \frac{z_1}{z_2} = z_1 z_2^{-1}$$

#### 7. Modulus or Absolute Value

$$|z| = \sqrt{x^2 + y^2}$$

#### 8. Conjugates and properties

$$\begin{aligned} \bar{z} &= x - iy = (x, -y) \\ \overline{z_1 \pm z_2} &= \bar{z}_1 \pm \bar{z}_2 \\ \overline{z_1 z_2} &= \bar{z}_1 \bar{z}_2 \\ \overline{\left(\frac{z_1}{z_2}\right)} &= \frac{\bar{z}_1}{\bar{z}_2} \end{aligned}$$

#### 9.

$$\begin{aligned} |z|^2 &= z\bar{z} \\ \operatorname{Re} z &= \frac{z + \bar{z}}{2}, \operatorname{Im} z = \frac{z - \bar{z}}{2i} \end{aligned}$$

#### 10. Triangle Inequality

$$|z_1 + z_2| \leq |z_1| + |z_2|$$

## Polar Coordinates and Euler Formula

#### 1. Polar Form: for $z \neq 0$ ,

$$z = r(\cos \theta + i \sin \theta)$$

where  $r = |z|$  and  $\tan \theta = y/x$ .  $\theta$  is called the argument of  $z$ . Since  $\theta + 2n\pi$  is also an argument of  $z$ , the principle value of argument of  $z$  is take such that  $-\pi < \theta \leq \pi$ . For  $z = 0$  the arg  $z$  is unde• ned.

#### 2. Euler formula: Symbollically,

$$e^{i\theta} = (\cos \theta + i \sin \theta)$$

#### 3.

$$\begin{aligned} z_1 z_2 &= r_1 r_2 e^{i(\theta_1 + \theta_2)} \\ \frac{z_1}{z_2} &= \frac{r_1}{r_2} e^{i(\theta_1 - \theta_2)} \\ z^n &= r^n e^{in\theta} \end{aligned}$$

#### 4. de Moivre's Formula

$$(\cos \theta + i \sin \theta)^n = \cos n\theta + i \sin n\theta$$



**Centurion**  
**UNIVERSITY**

Shaping Lives...  
Empowering Communities...

## Roots of Complex Numbers

Let  $z = r e^{i\theta}$  then

$$z^{1/n} = r^{1/n} \exp \left( i \left( \frac{\theta}{n} + \frac{2k\pi}{n} \right) \right)$$

There are only  $n$  distinct roots which can be given by  $k = 0, 1, \dots, n - 1$ . If  $\theta$  is a principle value of  $\arg z$  then  $\theta/n$  is called the principle root.

**Example 1.1** The three possible roots of  $\left(\frac{1+i}{\sqrt{2}}\right)^{1/3} = (e^{i\pi/4})^{1/3}$  are  $e^{i\pi/12}$ ,  $e^{i\pi/12+i2\pi/3}$ ,  $e^{i\pi/12+i4\pi/3}$ .

## Regions in Complex Plane

1.  $\epsilon$  - nbd of  $z_0$  is defined as a set of all points  $z$  which satisfy

$$|z - z_0| < \epsilon$$

2. Deleted nbd of  $z_0$  is a nbd of  $z_0$  excluding point  $z_0$ .

3. Interior Point, Exterior Point, Boundary Point, Open set and closed set.

4. Domain, Region, Bounded sets, Limit Points.



**Centurion**  
**UNIVERSITY**

Shaping Lives...  
Empowering Communities...

## Limits

A function  $f$  is defined in a deleted nbd of  $z_0$ .

**Definition 3.1** The limit of the function  $f(z)$  as  $z \rightarrow z_0$  is a number  $w_0$  if, for any given  $\epsilon > 0$  there exists a  $\delta > 0$  such that

$$|z - z_0| < \delta \Rightarrow |f(z) - w_0| < \epsilon.$$

**Example 3.1**  $f(z) = 5z$ . Show that  $\lim_{z \rightarrow z_0} f(z) = 5z_0$ .

**Example 3.2**  $f(z) = z^2$ . Show that  $\lim_{z \rightarrow z_0} f(z) = z_0^2$ .

**Example 3.3**  $f(z) = z/\bar{z}$ . Show that the limit of  $f$  does not exist as  $z \rightarrow 0$ .

**Theorem 3.1** Let  $f(z) = u(x, y) + iv(x, y)$  and  $w_0 = u_0 + iv_0$ .  $\lim_{z \rightarrow z_0} f(z) = w_0$  if and only if  $\lim_{(x,y) \rightarrow (x_0,y_0)} u = u_0$  and  $\lim_{(x,y) \rightarrow (x_0,y_0)} v = v_0$ .

**Example 3.4**  $f(z) = \sin z$ . Show that the  $\lim_{z \rightarrow z_0} f(z) = \sin z_0$

**Example 3.5**  $f(z) = 2x + iy^2$ . Show that the  $\lim_{z \rightarrow 2i} f(z) = 4i$ .

**Theorem 3.2** If  $\lim_{z \rightarrow z_0} f(z) = w_0$  and  $\lim_{z \rightarrow z_0} F(z) = W_0$ ,

$$\lim_{z \rightarrow z_0} [f(z) + F(z)] = w_0 + W_0;$$

$$\lim_{z \rightarrow z_0} f(z) F(z) = w_0 W_0;$$

$$\lim_{z \rightarrow z_0} f(z) / F(z) = \frac{w_0}{W_0} \quad W_0 \neq 0.$$

This theorem immediately makes available the entire machinery and tools used for real analysis to be applied to complex analysis. The rules for finding limits then can be listed as follows:



Centurion  
UNIVERSITY

Shaping Lives...  
Empowering Communities...

1.  $\lim_{z \rightarrow z_0} c = c.$
2.  $\lim_{z \rightarrow z_0} z^n = z_0^n.$
3.  $\lim_{z \rightarrow z_0} P(z) = P(z_0)$  if  $P$  is a polynomial in  $z.$
4.  $\lim_{z \rightarrow z_0} \exp(z) = \exp(z_0).$
5.  $\lim_{z \rightarrow z_0} \sin(z) = \sin z_0.$

## Continuity

**Definition 3.2** A function  $f$ , defined in some nbd of  $z_0$  is continuous at  $z_0$  if

$$\lim_{z \rightarrow z_0} f(z) = f(z_0).$$

This definition clearly assumes that the function is defined at  $z_0$  and the limit on the LHS exists. The function  $f$  is continuous in a region if it is continuous at all points in that region.

If functions  $f$  and  $g$  are continuous at  $z_0$  then  $f + g$ ,  $fg$  and  $f/g$  ( $g(z_0) \neq 0$ ) are also continuous at  $z_0$ .

If a function  $f(z) = u(x, y) + iv(x, y)$  is continuous at  $z_0$  then the component functions  $u$  and  $v$  are also continuous at  $(x_0, y_0)$ .

## Derivative

**Definition 3.3** A function  $f$ , defined in some nbd of  $z_0$  is differentiable at  $z_0$  if

$$\lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0}$$

exists. The limit is called the derivative of  $f$  at  $z_0$  and is denoted by  $f'(z_0)$  or  $\frac{df}{dz}(z_0)$ .

**Example 3.6**  $f(z) = z^2$ . Show that  $f'(z) = 2z$ .

**Example 3.7**  $f(z) = |z|^2$ . Show that this function is differentiable only at  $z = 0$ . In real analysis  $|x|$  is not differentiable but  $|x|^2$  is.

If a function is differentiable at  $z$ , then it is continuous at  $z$ .



Centurion  
UNIVERSITY

Shaping Lives...  
Empowering Communities

## Analytic Functions

**Definition 3.4** A function is analytic in an open set if it has a derivative at each point in that set.

**Definition 3.5** A function is analytic at a point  $z_0$  if it is analytic in some nbd of  $z_0$ .

**Definition 3.6** A function is an entire function if it is analytic at all points of  $C$ .

**Example 3.13**  $f(z) = 1/z$  is analytic at all nonzero points.

**Example 3.14**  $f(z) = |z|^2$  is not analytic anywhere.

A function is not analytic at a point  $z_0$ , but is analytic at some point in each nbd of  $z_0$  then  $z_0$  is called the singular point of the function  $f$ .

## Harmonic Functions

**Definition 3.7** A real valued function  $H(x, y)$  is said to be harmonic in a domain of  $xy$  plane if it has continuous partial derivatives of the first and second order and satisfies Laplace equation:

$$H_{xx}(x, y) + H_{yy}(x, y) = 0.$$

# Cauchy-Riemann Equations



**Centurion**  
**UNIVERSITY**

Shaping Lives...  
Empowering Communities...

**Theorem 3.3** If  $f'(z_0)$  exists, then all the first order partial derivatives of component function  $u(x, y)$  and  $v(x, y)$  exist and satisfy Cauchy-Riemann Conditions:

$$u_x = v_y$$

$$u_y = -v_x.$$

**Example 3.8**  $f(z) = z^2 = x^2 - y^2 + i2xy$ . Show that Cauchy-Riemann Conditions are satisfied.

**Example 3.9**  $f(z) = |z|^2 = x^2 + y^2$ . Show that the Cauchy-Riemann Conditions are satisfied only at  $z = 0$ .

**Theorem 3.4** Let  $f(z) = u(x, y) + iv(x, y)$  be defined in some neighborhood of the point  $z_0$ . If the first partial derivatives of  $u$  and  $v$  exist and are continuous at  $z_0$  and satisfy Cauchy-Riemann equations at  $z_0$ , then  $f$  is differentiable at  $z_0$  and

$$f'(z) = u_x + iv_x = v_y - iu_y.$$

**Example 3.10**  $f(z) = \exp(z)$ . Show that  $f'(z) = \exp(z)$ .

**Example 3.11**  $f(z) = \sin(z)$ . Show that  $f'(z) = \cos(z)$ .

**Example 3.12**  $f(z) = \frac{z^2}{z}$ . Show that the CR conditions are satisfied at  $z = 0$  but the function  $f$  is not differentiable at 0.



Centurion  
UNIVERSITY

Shaping Lives...  
Empowering Communities...

1. The *Cauchy-Riemann equations* for two functions  $u(x, y)$  and  $v(x, y)$  are

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \quad \text{and} \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$$

- (a) Show that  $u(x, y) = x^2 - y^2$  and  $v(x, y) = 2xy$  satisfy the Cauchy-Riemann equations.
- (b) The *Laplace equation* for a function  $f(x, y)$  is

$$f_{xx} + f_{yy} = 0.$$

Suppose that  $a$  and  $b$  satisfy the Cauchy-Riemann equations and have equal mixed partial derivatives (i.e.  $a_{xy} = a_{yx}$ , etc.).

Show that  $a + b$  satisfies the Laplace equation.




**Centurion**  
**UNIVERSITY**

Shaping Lives...  
Empowering Communities...

# Cauchy Integral Formula


**R**- a simply connected region



#.  $f(z)$  is assumed analytic in  $R$  but we multiply by a factor  $\frac{1}{(z - z_0)}$  that is analytic except at  $z_0$  and consider the integral around  $C$

$$\int_C \frac{f(z)}{(z - z_0)} dz$$

#. To evaluate, consider the path  $C + c_1 + c_2 + C_0$  shown that encloses a simply connected region for which the integrand is analytic on and inside the path:



$$\int_{C + c_1 + c_2 + C_0} \frac{f(z)}{(z - z_0)} dz = 0 \Rightarrow \int_C \frac{f(z)}{(z - z_0)} dz = - \int_{C_0} \frac{f(z)}{(z - z_0)} dz$$



**Centurion**  
**UNIVERSITY**

Shaping Lives...  
Empowering Communities...

# Cauchy Integral Formula, cont'd

$$\int_C \frac{f(z)}{z-z_0} dz = - \int_{C_0} \frac{f(z)}{z-z_0} dz$$



Evaluate the  $C_0$  integral on a circular path,  $z-z_0 = re^{i\theta}$ ,  $dz = rie^{i\theta} d\theta$

$$\int_{C_0} \frac{f(z)}{(z-z_0)} dz \stackrel{r \rightarrow 0}{=} f(z_0) \int_{2\pi}^0 \frac{f(z_0) i e^{i\theta} d\theta}{f(z_0) e^{i\theta}} = -2\pi i f(z_0)$$

for  $r \rightarrow 0$

$$\Rightarrow \int_C \frac{f(z)}{z-z_0} dz = 2\pi i f(z_0)$$

$$\Rightarrow f(z_0) = \frac{1}{2\pi i} \oint_C \frac{f(z)}{z-z_0} dz$$

Cauchy Integral  
Formula


The value of  $f(z)$  at  $z_0$  is completely determined by its values on  $C$ !



Centurion  
UNIVERSITY

Shaping Lives...  
Empowering Communities...

# Cauchy Integral Formula, cont'd



Note that if  $z_0$  is outside  $C$ , the integrand is analytic inside  $C$ , hence by the Cauchy Integral Theorem,

$$\frac{1}{2\pi i} \oint_C \frac{f(z)}{z - z_0} dz = 0$$

In summary,

$$\oint_C \frac{f(z)}{z - z_0} dz = \begin{cases} 2\pi i f(z_0), & z_0 \text{ inside } C \\ 0, & z_0 \text{ outside } C \end{cases}$$



**Centurion**  
**UNIVERSITY**

Shaping Lives...  
Empowering Communities...

## Contour Integral

If  $C$  is a contour in complex plane defined by  $z(t) = x(t) + iy(t)$  and a function  $f(z) = u(x, y) + iv(x, y)$  is defined on it. The integral of  $f(z)$  along the contour  $C$  is denoted and defined as follows:

$$\begin{aligned} \int_C f(z) dz &= \int_a^b f(z) z'(t) dt \\ &= \int_a^b (ux' - vy') dt + i \int_a^b (uy' + vx') dt \\ &= \int (udx - vdy) + i \int (udy + vdx) \end{aligned}$$

The component integrals are usual real integrals and are well defined. In the last form appropriate limits must be placed in the integrals.

Some very straightforward rules of integration are given below:

1.  $\int_C w f(z) dz = w \int_C f(z) dz$  where  $w$  is a complex constant.
2.  $\int_C (f(z) + g(z)) dz = \int_C f(z) dz + \int_C g(z) dz$ .
3.  $\int_{C_1+C_2} f(z) dz = \int_{C_1} f(z) dz + \int_{C_2} f(z) dz$ .
4.  $|\int_C f(z) dz| \leq \int_C |f(z(t)) z'(t)| dt$ .
5. If  $|f(z)| \leq M$  for all  $z \in C$  then  $|\int_C f(z) dz| \leq ML$ , where  $L$  is length of the contour  $C$ .

**Example 4.4**  $f(z) = z^2$ . Find integral of  $f$  from  $(0, 0)$  to  $(2, 1)$  along a straight line and also along st line path from  $(0, 0)$  to  $(2, 0)$  and from  $(2, 0)$  to  $(2, 1)$ .

**Example 4.5**  $f(z) = 1/z$ . Find the integral from  $(2, 0)$  to  $(-2, 0)$  along a semicircular path in upper plane given by  $|z| = 2$ .

**Example 4.6** Show that  $|\int_C f(z) dz| \leq \frac{\pi}{3}$  for  $f(z) = 1/(z^2 - 1)$  and  $C : |z| = 2$  from  $2$  to  $2i$ .



**Centurion**  
**UNIVERSITY**

Shaping Lives...  
Empowering Communities...

## Cauchy-Goursat Theorem

**Theorem 4.1 (Jordan Curve Theorem)** Every simple and closed contour in complex plane splits the entire plane into two domains one of which is bounded. The bounded domain is called the interior of the contour and the other one is called the exterior of the contour.

Define a sense direction for a contour.

**Theorem 4.2** Let  $C$  be a simple closed contour with positive orientation and let  $D$  be the interior of  $C$ . If  $P$  and  $Q$  are continuous and have continuous partial derivatives  $P_x, P_y, Q_x$  and  $Q_y$  at all points on  $C$  and  $D$ , then

$$\int_C (P(x, y)dx + Q(x, y)dy) = \iint_D [Q_x(x, y) - P_y(x, y)] dx dy$$

**Theorem 4.3 (Cauchy-Goursat Theorem)** Let  $f$  be analytic in a simply connected domain  $D$ . If  $C$  is any simple closed contour in  $D$ , then

$$\int_C f(z) dz = 0.$$

**Example 4.7**  $f(z) = z^2, \exp(z), \cos(z)$  etc are entire functions so integral about any loop is zero.

**Theorem 4.4** Let  $C_1$  and  $C_2$  be two simple closed positively oriented contours such that  $C_2$  lies entirely in the interior of  $C_1$ . If  $f$  is an analytic function in a domain  $D$  that contains  $C_1$  and  $C_2$  both and the region between them, then

$$\int_{C_1} f(z) dz = \int_{C_2} f(z) dz.$$

**Example 4.8**  $f(z) = 1/z$ . Find  $\int_C f(z) dz$  if  $C$  is any contour containing origin. Choose a circular contour inside  $C$ .

**Example 4.9**  $\int_C \frac{1}{z-z_0} dz = 2\pi i$  if  $C$  contains  $z_0$ .

**Example 4.10** Find  $\int_C \frac{2z dz}{z^2+2}$  where  $C : |z| = 2$ . Extend the Cauchy Goursat theorem to multiply connected domains.



**Centurion**  
**UNIVERSITY**

Shaping Lives...  
Empowering Communities...

## Singularities

**Definition 6.1** *If a function  $f$  fails to be analytic at  $z_0$  but is analytic at some point in each neighbourhood of  $z_0$ , then  $z_0$  is a singular point of  $f$ .*

**Definition 6.2** *If a function  $f$  fails to be analytic at  $z_0$  but is analytic at each  $z$  in  $0 < |z - z_0| < \delta$  for some  $\delta$ , then  $f$  is said to be an isolated singular point of  $f$ .*

**Example 6.1**  $f(z) = 1/z$  has an isolated singularity at 0.

**Example 6.2**  $f(z) = 1/\sin(\pi z)$  has isolated singularities at  $z = 0, \pm 1, \dots$

**Example 6.3**  $f(z) = 1/\sin(\pi/z)$  has isolated singularities at  $z = 1/n$  for integral  $n$ , also has a singularity at  $z = 0$ .

**Example 6.4**  $f(z) = \log z$  all points of negative  $x$ -axis are singular.

## Types of singularities

If a function  $f$  has an isolated singularity at  $z_0$  then  $\exists$  a  $\delta$  such that  $f$  is analytic at all points in  $0 < |z - z_0| < \delta$ . Then  $f$  must have a Laurent series expansion about  $z_0$ . The part  $\sum_{n=1}^{\infty} b_n (z - z_0)^{-n}$  is called the principal part of  $f$ .

1. If there are infinite nonzero  $b_i$  in the principal part then  $z_0$  is called an essential singularity of  $f$ .
2. If for some integer  $m$ ,  $b_m \neq 0$  but  $b_i = 0$  for all  $i > m$  then  $z_0$  is called a pole of order  $m$  of  $f$ . If  $m = 1$  then it is called a simple pole.
3. If all  $b$ s are zero then  $z_0$  is called a removable singularity.



**Centurion**  
**UNIVERSITY**

Shaping Lives...  
Empowering Communities...

## Residues

Suppose a function  $f$  has an isolated singularity at  $z_0$ , then there exists a  $\delta > 0$  such that  $f$  is analytic for all  $z$  in deleted nbd  $0 < |z - z_0| < \delta$ . Then  $f$  has a Laurent series representation

$$f(z) = \sum_{n=0}^{\infty} a_n (z - z_0)^n + \sum_{n=1}^{\infty} \frac{b_n}{(z - z_0)^n}.$$

The coefficient

$$b_1 = \frac{1}{2\pi i} \int_C f(z) dz$$

where  $C$  is any contour in the deleted nbd, is called the residue of  $f$  at  $z_0$ .

**Example 6.6**  $f(z) = \frac{1}{z-z_0}$ . Then  $\int_C f(z) dz = 2\pi i b_1 = 2\pi i$  if  $C$  contains  $z_0$ , otherwise 0.

**Example 6.7**  $f(z) = \frac{1}{z(z-z_0)^2}$ . Show  $\int_C f(z) dz = -\frac{\pi i}{8}$  if  $C : |z - 2| = 1$ .

**Example 6.8**  $f(z) = z \exp\left(\frac{1}{z}\right)$ . Show  $\int_C f(z) dz = \pi i$  if  $C : |z| = 1$ .

**Example 6.9**  $f(z) = \exp\left(\frac{1}{z^2}\right)$ . Show  $\int_C f(z) dz = 0$  if  $C : |z| = 1$  even though it has a singularity at  $z = 0$ .

**Theorem 6.1** If a function  $f$  is analytic on and inside a positively oriented contour  $C$ , except for a finite number of points  $z_1, z_2, \dots, z_k$  inside  $C$ , then

$$\int_C f(z) dz = 2\pi i \sum_{i=1}^k \text{Res} f(z_i).$$

**Example 6.10** Show that  $\int \frac{5z-2}{z(z-1)} dz = 10\pi i$

## Residues of Poles

**Theorem 6.2** If a function  $f$  has a pole of order  $m$  at  $z_0$  then

$$\text{Res} f(z_0) = \lim_{z \rightarrow z_0} \left[ \frac{1}{(m-1)!} \frac{d^{m-1}}{dz^{m-1}} ((z - z_0)^m f(z)) \right].$$

**Example 6.11**  $f(z) = \frac{1}{z-z_0}$ . Simple pole  $\text{Res} f(z_0) = 1$ .



**Centurion**  
**UNIVERSITY**

Shaping Lives...  
Empowering Communities...

**Example 6.12**  $f(z) = \frac{1}{z(z-2)^4}$ . Simple pole at  $z = 0$ .  $\text{Res } f(0) = 1/16$ . Pole of order 4 at  $z = 2$ .  $\text{Res } f(2) = -1/16$ .

**Example 6.13**  $f(z) = \frac{\cos z}{z^2(z-\pi)^3}$ .  $\text{Res } f(0) = -3/\pi^4$ .  $\text{Res } f(\pi) = -(6 - \pi^2)/2\pi^4$ .

## Quotients of Analytic Functions

**Theorem 6.3** If a function  $f(z) = \frac{P(z)}{Q(z)}$ , where  $P$  and  $Q$  are analytic at  $z_0$ , then

1.  $f$  is singular at  $z_0$  iff  $Q(z_0) = 0$ .
2.  $f$  has a simple pole at  $z_0$  if  $Q'(z_0) \neq 0$ . Then residue of  $f$  at  $z_0$  is  $P(z_0)/Q'(z_0)$ .




Centurion  
UNIVERSITY

Shaping Lives...  
Empowering Communities...

# Singularities

## Isolated singularities



$$\frac{\sin(z)}{z}, \frac{1 - \cos(z)}{z^2}$$

$$\frac{1}{z}, \frac{1}{z^2}, \frac{1}{(z-1)^m},$$

$$\frac{2z+3}{(z-1)^2(z+2)}$$

$$\sin\left(\frac{1}{z}\right), e^{1/z}$$

These are each discussed in more detail next.



**Centurion**  
**UNIVERSITY**

Shaping Lives...  
Empowering Communities...

Essential singularities can be distinguished from poles by the fact that they cannot be removed by multiplying by a factor of finite value.

Example:

$$w = e^{1/z} = 1 + \frac{1}{z} + \frac{1}{2! z^2} + \dots + \frac{1}{n! z^n} + \dots \leftarrow \text{infinite at the origin}$$

We try to remove the singularity of the function at the origin by multiplying  $z^p$

$$z^p w = z^p + z^{p-1} + \frac{z^{p-2}}{2!} + \dots + \frac{z^{p-n}}{n!} + \dots$$

It consists of a finite number of positive powers of  $z$ , followed by an infinite number of negative powers of  $z$ .

All terms are positive

$$\text{As } z \rightarrow 0, \quad z^p w \rightarrow \infty$$

It is impossible to find a finite value of  $p$  which will remove the singularity in  $e^{1/z}$  at the origin.

The singularity is “essential”.



Centurion  
UNIVERSITY

Shaping Lives...  
Empowering Communities...

# Singularities

## Examples of singularities:

(These will be discussed in more detail later.)

T  $\frac{\sin(z)}{z}$  removable singularity at  $z = 0$

L  $\frac{1}{(z - z_s)^p}$  pole of order  $p$  at  $z = z_s$  ( if  $p = 1$ , pole is a *simple pole*)

L  $e^{1/z}$  isolated essential singularity at  $z = z_0$  (pole of infinite order)

N  $\frac{1}{\sin\left(\frac{1}{z}\right)}$  non-isolated essential singularity  $z = 0$

N  $z^{1/2}$  branch point (*not* an isolated singularity)

If expanded about the singularity, we can have:

T = Taylor

L = Laurent

N = Neither



Centurion  
UNIVERSITY


Shaping Lives...  
Empowering Communities...

# Barrier - Branch Cut

The barrier must start from the branch point but it can go to infinity in any direction in the  $z$  plane, and may be either curved or straight.

In most normal applications, the barrier is drawn along the negative real axis.

- The branch is termed the “principle branch”.
- The barrier is termed the “branch cut”.
- For the example given in the previous slide, the region, the barrier confines the function to the region in which the argument of  $z$  is within the range  $-\pi < \theta < \pi$ .






Centurion  
UNIVERSITY

Shaping Lives...  
Empowering Communities...

# Branch points

- The singularities described above arise from the non-analytic behaviour of single-valued functions.
- However, multi-valued functions frequently arise in the solution of engineering problems.
- For example:

$$\boxed{w = z^{\frac{1}{2}}} \xrightarrow{z = re^{i\theta}} \boxed{w = r^{\frac{1}{2}} e^{\frac{1}{2}i\theta}}$$



For any value of  $z$  represented by a point on the circumference of the circle in the  $z$  plane, there will be two corresponding values of  $w$  represented by points in the  $w$  plane.



**Centurion**  
**UNIVERSITY**

Shaping Lives...  
Empowering Communities...

# Laurent's Theorem

→ **Laurent series** of a complex function  $f(z)$  is a representation of that function as a power series which includes terms of negative degree. It may be used to express complex functions in cases where a Taylor series expansion cannot be applied.


$$f(z) = \sum_{n=-\infty}^{\infty} a_n (z - c)^n$$

$$\begin{aligned} C &= z_0 \\ K &= n \\ A &= z_0 \end{aligned}$$

Also say that ,

$$f(z) = \sum_{k=0}^{\infty} a_k (z - a)^k + \sum_{k=1}^{\infty} b_k (z - a)^{-k},$$

$$a_n = \frac{1}{2\pi i} \oint_{\gamma} \frac{f(z) dz}{(z - c)^{n+1}}$$





Centurion  
UNIVERSITY

Shaping Lives...  
Empowering Communities...

# Taylor's series

- **Taylor series** is a representation of a function as an infinite sum of terms that are calculated from the values of the function's derivatives at a single point.

$$f(x) =$$

$$f(a) + f'(a)(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \frac{f^{(3)}(a)}{3!}(x-a)^3 + \dots + \frac{f^{(n)}(a)}{n!}(x-a)^n + \dots$$

Here,  $a = z_0$ , center  
 $x = z$  inside circle  $C$



**Centurion  
UNIVERSITY**

Shaping Lives...  
Empowering Communities...


# The Residue Theorem

Section 8

We saw that the integral of an analytic function  $f(z)$  over a closed curve  $C$  when  $f(z)$  has one singular point inside  $C$  is

$$\oint_C f(z) dz = 2\pi j b_1$$


$b_1$  is the residue of  $f(z)$  at  $z_0$



Residue Theorem: Let  $f(z)$  be an analytic function inside and on a closed path  $C$  except for at  $k$  singular points inside  $C$ .

Then

$$\oint_C f(z) dz = 2\pi j \sum_{i=1}^k \text{Res}_{z=z_i} f(z)$$





**Centurion**  
**UNIVERSITY**

*Shaping Lives...*  
*Empowering Communities...*

Numericals from various sources



**Centurion**  
**UNIVERSITY**

*Shaping Lives...*  
*Empowering Communities...*

# **Module 4 , 5 , 6 , 7**

## **Integral Transforms**



# Fourier Integral

If  $f(x)$  and  $f'(x)$  are piecewise continuous in every finite interval, and  $f(x)$  is absolutely integrable on  $\mathbb{R}$ , i.e.

converges, then

$$\frac{1}{2} [f(x-) + f(x+)] = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{iwx} \left( \int_{-\infty}^{\infty} e^{-iwt} f(t) dt \right) dw$$

Remark: the above conditions are sufficient, but not necessary.



Centurion  
UNIVERSITY

Shaping Lives...  
Empowering Communities...

$$\frac{y}{\sqrt{c}} = \sqrt{\frac{2}{\pi}} \int_0^{\infty} f(x) \cos kx \, dx = F_c(k)$$

} Fourier cosine  
transform

$$\frac{y}{\sqrt{c}}^{-1} = \sqrt{\frac{2}{\pi}} \int_0^{\infty} F_c(k) \cos kx \, dk = f(x)$$

Fourier sine Transform:

$$\frac{y}{\sqrt{s}} = \sqrt{\frac{2}{\pi}} \int_0^{\infty} f(x) \sin kx \, dx = F_s(k)$$

} Fourier sine  
transform

$$\frac{y}{\sqrt{s}}^{-1} = \sqrt{\frac{2}{\pi}} \int_0^{\infty} F_s(k) \sin kx \, dk = f(x)$$



**Centurion  
UNIVERSITY**

Shaping Lives...  
Empowering Communities...

► Find the Fourier transform of the normalised Gaussian distribution

$$f(t) = \frac{1}{\tau\sqrt{2\pi}} \exp\left(-\frac{t^2}{2\tau^2}\right), \quad -\infty < t < \infty.$$

This Gaussian distribution is centred on  $t = 0$  and has a root mean square deviation  $\Delta t = \tau$ . (Any reader who is unfamiliar with this interpretation of the distribution should refer to chapter 30.)

Using the definition (13.5), the Fourier transform of  $f(t)$  is given by

$$\begin{aligned} \tilde{f}(\omega) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{1}{\tau\sqrt{2\pi}} \exp\left(-\frac{t^2}{2\tau^2}\right) \exp(-i\omega t) dt \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{1}{\tau\sqrt{2\pi}} \exp\left\{-\frac{1}{2\tau^2} [t^2 + 2\tau^2 i\omega t + (\tau^2 i\omega)^2 - (\tau^2 i\omega)^2]\right\} dt, \end{aligned}$$

where the quantity  $-(\tau^2 i\omega)^2/(2\tau^2)$  has been both added and subtracted in the exponent in order to allow the factors involving the variable of integration  $t$  to be expressed as a complete square. Hence the expression can be written

$$\tilde{f}(\omega) = \frac{\exp(-\frac{1}{2}\tau^2\omega^2)}{\sqrt{2\pi}} \left\{ \frac{1}{\tau\sqrt{2\pi}} \int_{-\infty}^{\infty} \exp\left[-\frac{(t + i\tau^2\omega)^2}{2\tau^2}\right] dt \right\}.$$

The quantity inside the braces is the normalisation integral for the Gaussian and equals unity, although to show this strictly needs results from complex variable theory (chapter 24). That it is equal to unity can be made plausible by changing the variable to  $s = t + i\tau^2\omega$  and assuming that the imaginary parts introduced into the integration path and limits (where the integrand goes rapidly to zero anyway) make no difference.

We are left with the result that

$$\tilde{f}(\omega) = \frac{1}{\sqrt{2\pi}} \exp\left(\frac{-\tau^2\omega^2}{2}\right), \quad (13.7)$$

which is another Gaussian distribution, centred on zero and with a root mean square deviation  $\Delta\omega = 1/\tau$ . It is interesting to note, and an important property, that the Fourier transform of a Gaussian is another Gaussian. ◀

In the above example the root mean square deviation in  $t$  was  $\tau$ , and so it is seen that the deviations or 'spreads' in  $t$  and in  $\omega$  are inversely related:

$$\Delta\omega \Delta t = 1,$$

independently of the value of  $\tau$ . In physical terms, the narrower in time is, say, an electrical impulse the greater the spread of frequency components it must contain. Similar physical statements are valid for other pairs of Fourier-related variables, such as spatial position and wave number. In an obvious notation,  $\Delta k \Delta x = 1$  for a Gaussian wave packet.

The uncertainty relations as usually expressed in quantum mechanics can be related to this if the de Broglie and Einstein relationships for momentum and energy are introduced; they are

$$p = \hbar k \quad \text{and} \quad E = \hbar\omega.$$



**Centurion**  
**UNIVERSITY**

Shaping Lives...  
Empowering Communities...

### 13.1.3 The Dirac $\delta$ -function

Before going on to consider further properties of Fourier transforms we make a digression to discuss the Dirac  $\delta$ -function and its relation to Fourier transforms. The  $\delta$ -function is different from most functions encountered in the physical sciences but we will see that a rigorous mathematical definition exists; the utility of the  $\delta$ -function will be demonstrated throughout the remainder of this chapter. It can be visualised as a very sharp narrow pulse (in space, time, density, etc.) which produces an integrated effect having a definite magnitude. The formal properties of the  $\delta$ -function may be summarised as follows.

The Dirac  $\delta$ -function has the property that

$$\delta(t) = 0 \quad \text{for } t \neq 0, \quad (13.11)$$

but its fundamental defining property is

$$\int f(t)\delta(t - a) dt = f(a), \quad (13.12)$$

provided the range of integration includes the point  $t = a$ ; otherwise the integral



**Centurion**  
**UNIVERSITY**

Shaping Lives...  
Empowering Communities...

equals zero. This leads immediately to two further useful results:

$$\int_{-a}^b \delta(t) dt = 1 \quad \text{for all } a, b > 0 \quad (13.13)$$

and

$$\int \delta(t - a) dt = 1, \quad (13.14)$$

provided the range of integration includes  $t = a$ .

Equation (13.12) can be used to derive further useful properties of the Dirac  $\delta$ -function:

$$\delta(t) = \delta(-t), \quad (13.15)$$

$$\delta(at) = \frac{1}{|a|} \delta(t), \quad (13.16)$$

$$t\delta(t) = 0. \quad (13.17)$$



Centurion  
UNIVERSITY

Shaping Lives...  
Empowering Communities...

► Prove that  $\delta(bt) = \delta(t)/|b|$ .

Let us first consider the case where  $b > 0$ . It follows that

$$\int_{-\infty}^{\infty} f(t)\delta(bt) dt = \int_{-\infty}^{\infty} f\left(\frac{t'}{b}\right) \delta(t') \frac{dt'}{b} = \frac{1}{b} f(0) = \frac{1}{b} \int_{-\infty}^{\infty} f(t)\delta(t) dt,$$

where we have made the substitution  $t' = bt$ . But  $f(t)$  is arbitrary and so we immediately see that  $\delta(bt) = \delta(t)/b = \delta(t)/|b|$  for  $b > 0$ .

Now consider the case where  $b = -c < 0$ . It follows that

$$\begin{aligned} \int_{-\infty}^{\infty} f(t)\delta(bt) dt &= \int_{\infty}^{-\infty} f\left(\frac{t'}{-c}\right) \delta(t') \left(\frac{dt'}{-c}\right) = \int_{-\infty}^{\infty} \frac{1}{c} f\left(\frac{t'}{-c}\right) \delta(t') dt' \\ &= \frac{1}{c} f(0) = \frac{1}{|b|} f(0) = \frac{1}{|b|} \int_{-\infty}^{\infty} f(t)\delta(t) dt, \end{aligned}$$

where we have made the substitution  $t' = bt = -ct$ . But  $f(t)$  is arbitrary and so

$$\delta(bt) = \frac{1}{|b|} \delta(t),$$

for all  $b$ , which establishes the result. ◀



Centurion  
UNIVERSITY

Shaping Lives...  
Empowering Communities...

(i) Differentiation:

$$\mathcal{F}[f'(t)] = i\omega\tilde{f}(\omega). \quad (13.28)$$

This may be extended to higher derivatives, so that

$$\mathcal{F}[f''(t)] = i\omega\mathcal{F}[f'(t)] = -\omega^2\tilde{f}(\omega),$$

and so on.

(ii) Integration:

$$\mathcal{F}\left[\int^t f(s) ds\right] = \frac{1}{i\omega}\tilde{f}(\omega) + 2\pi c\delta(\omega), \quad (13.29)$$

where the term  $2\pi c\delta(\omega)$  represents the Fourier transform of the constant of integration associated with the indefinite integral.

(iii) Scaling:

$$\mathcal{F}[f(at)] = \frac{1}{a}\tilde{f}\left(\frac{\omega}{a}\right). \quad (13.30)$$

(iv) Translation:

$$\mathcal{F}[f(t+a)] = e^{ia\omega}\tilde{f}(\omega). \quad (13.31)$$

(v) Exponential multiplication:

$$\mathcal{F}[e^{\alpha t}f(t)] = \tilde{f}(\omega + i\alpha), \quad (13.32)$$

where  $\alpha$  may be real, imaginary or complex.



**Centurion**  
**UNIVERSITY**

Shaping Lives...  
Empowering Communities...

► Prove relation (13.28).

Calculating the Fourier transform of  $f'(t)$  directly, we obtain

$$\begin{aligned}\mathcal{F}[f'(t)] &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f'(t) e^{-i\omega t} dt \\ &= \frac{1}{\sqrt{2\pi}} \left[ e^{-i\omega t} f(t) \right]_{-\infty}^{\infty} + \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} i\omega e^{-i\omega t} f(t) dt \\ &= i\omega \tilde{f}(\omega),\end{aligned}$$

if  $f(t) \rightarrow 0$  at  $t = \pm\infty$ , as it must since  $\int_{-\infty}^{\infty} |f(t)| dt$  is finite. ◀

## Example 15.4.2 HEAT FLOW PDE



Centurion  
UNIVERSITY

Shaping Lives...  
Empowering Communities...

To illustrate another transformation of a PDE into an ODE, let us Fourier transform the heat flow partial differential equation

$$\frac{\partial \psi}{\partial t} = a^2 \frac{\partial^2 \psi}{\partial x^2},$$

where the solution  $\psi(x, t)$  is the temperature in space as a function of time. By taking the Fourier transform of both sides of this equation (note that here only  $\omega$  is the transform variable conjugate to  $x$  because  $t$  is the time in the heat flow PDE), where

$$\Psi(\omega, t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \psi(x, t) e^{i\omega x} dx,$$

this yields an ODE for the Fourier transform  $\Psi$  of  $\psi$  in the time variable  $t$ ,

$$\frac{\partial \Psi(\omega, t)}{\partial t} = -a^2 \omega^2 \Psi(\omega, t).$$

Integrating we obtain

$$\ln \Psi = -a^2 \omega^2 t + \ln C, \quad \text{or} \quad \Psi = C e^{-a^2 \omega^2 t},$$

where the integration constant  $C$  may still depend on  $\omega$  and, in general, is determined by initial conditions. In fact,  $C = \Psi(\omega, 0)$  is the initial spatial distribution of  $\Psi$ , so it is given by the transform (in  $x$ ) of the initial distribution of  $\psi$ , namely,  $\psi(x, 0)$ . Putting this solution back into our inverse Fourier transform, this yields

$$\psi(x, t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} C(\omega) e^{-i\omega x} e^{-a^2 \omega^2 t} d\omega.$$

For simplicity, we here take  $C$   $\omega$ -independent (assuming a delta-function initial temperature distribution) and integrate by completing the square in  $\omega$ , as in Example 15.1.1,



**Centurion**  
**UNIVERSITY**

Shaping Lives...  
Empowering Communities...

making appropriate changes of variables and parameters ( $a^2 \rightarrow a^2 t$ ,  $\omega \rightarrow x$ ,  $t \rightarrow -\omega$ ). This yields the particular solution of the heat flow PDE,

$$\psi(x, t) = \frac{C}{a\sqrt{2t}} \exp\left(-\frac{x^2}{4a^2 t}\right),$$

which appears as a clever guess in Chapter 8. In effect, we have shown that  $\psi$  is the inverse Fourier transform of  $C \exp(-a^2 \omega^2 t)$ . ■

**Example 15.4.3** Inverse of PDE



**Centurion**  
**UNIVERSITY**

Shaping Lives...  
Empowering Communities...

## 15.5 CONVOLUTION THEOREM

We shall employ convolutions to solve differential equations, to normalize momentum wave functions (Section 15.6), and to investigate transfer functions (Section 15.7).

Let us consider two functions  $f(x)$  and  $g(x)$  with Fourier transforms  $F(t)$  and  $G(t)$ , respectively. We define the operation

$$f * g \equiv \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} g(y) f(x - y) dy \quad (15.52)$$

as the **convolution** of the two functions  $f$  and  $g$  over the interval  $(-\infty, \infty)$ . This form of an integral appears in probability theory in the determination of the probability density of two random, independent variables. Our solution of Poisson's equation, Eq. (9.148), may be interpreted as a convolution of a charge distribution,  $\rho(\mathbf{r}_2)$ , and a weighting function,  $(4\pi\epsilon_0|\mathbf{r}_1 - \mathbf{r}_2|)^{-1}$ . In other works this is sometimes referred to as the **Faltung**, to use the



Centurion  
UNIVERSITY

Shaping Lives...  
Empowering Communities...




FIGURE 15.5

German term for “folding.”<sup>5</sup> We now transform the integral in Eq. (15.52) by introducing the Fourier transforms:

$$\begin{aligned}
 \int_{-\infty}^{\infty} g(y) f(x-y) dy &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} g(y) \int_{-\infty}^{\infty} F(t) e^{-it(x-y)} dt dy \\
 &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} F(t) \left[ \int_{-\infty}^{\infty} g(y) e^{ity} dy \right] e^{-itx} dt \\
 &= \int_{-\infty}^{\infty} F(t) G(t) e^{-itx} dt, \tag{15.53}
 \end{aligned}$$

interchanging the order of integration and transforming  $g(y)$ . This result may be interpreted as follows: The Fourier inverse transform of a **product** of Fourier transforms is the convolution of the original functions,  $f * g$ .

For the special case  $x = 0$  we have

$$\int_{-\infty}^{\infty} F(t) G(t) dt = \int_{-\infty}^{\infty} f(-y) g(y) dy. \tag{15.54}$$

The minus sign in  $-y$  suggests that modifications be tried. We now do this with  $g^*$  instead of  $g$  using a different technique.



Centurion  
UNIVERSITY

Shaping Lives...  
Empowering Communities...

# The Laplace Transform

## Time Integration:

The property is:

$$L\left[\int_0^{\infty} f(t)dt\right] = \int_0^{\infty} \left[\int_0^t f(x)dx\right] e^{-st} dt$$

*Integrate by parts:*

$$\text{Let } u = \int_0^t f(x)dx, \quad du = f(t)dt$$

*and*

$$dv = e^{-st} dt, \quad v = -\frac{1}{s} e^{-st}$$



**Centurion**  
**UNIVERSITY**

*Shaping Lives...*  
*Empowering Communities...*

Home Task:

For Inverse Laplace  
transform material go  
through Arfken  
available in your library.



**Centurion**  
**UNIVERSITY**

Shaping Lives...  
Empowering Communities...

## *Real Integration Theorem*

This theorem establishes the relationship between the Laplace transform of a function and that of its integral. It states that

$$\mathcal{L} \left[ \int_0^t f(t) dt \right] = \frac{1}{s} F(s) \quad (2-1.7)$$

The proof of this theorem is carried out by integrating the definition of the Laplace transform by parts. This proof is similar to that of the real differentiation theorem and is left as an exercise. The Laplace transform of the  $n$ th integral of a function is the transform of the function divided by  $s^n$ .



**Centurion**  
**UNIVERSITY**

Shaping Lives...  
Empowering Communities...

## Laplace Transforms of Common Functions

Function	Laplace Transform	Restrictions
1	$\frac{1}{s}$	$s > 0$
$e^{at}$	$\frac{1}{s-a}$	$s > a$
$t^n$	$\frac{n!}{s^{n+1}}$	$s > 0, n$ an integer $> 0$
$\cos at$	$\frac{s}{s^2 + a^2}$	$s > 0$
$\sin at$	$\frac{a}{s^2 + a^2}$	$s > 0$
$\cosh at$	$\frac{s}{s^2 - a^2}$	$s >  a $
$\sinh at$	$\frac{a}{s^2 - a^2}$	$s >  a $
$e^{at} \cos bt$	$\frac{s-a}{(s-a)^2 + b^2}$	$s > a$
$e^{at} \sin bt$	$\frac{b}{(s-a)^2 + b^2}$	$s > a$
$t^n e^{at}$	$\frac{n!}{(s-a)^{n+1}}$	$s > a, n$ an integer $> 0$
$f(ct)$	$\frac{1}{c} \mathcal{L}\{f(s/c)\}$	$c > 0$
$f^{(n)}(t)$	$s^n \mathcal{L}\{f(t)\} - s^{n-1}f(0) - \dots - sf^{(n-2)}(0) - f^{(n-1)}(0)$	



**Centurion**  
**UNIVERSITY**

Shaping Lives...  
Empowering Communities...

## Table of Inverse Laplace Transform

$$1. \mathcal{L}^{-1} \left\{ \frac{1}{s} \right\} = 1$$

$$2. \mathcal{L}^{-1} \left\{ \frac{1}{s - a} \right\} = e^{at}$$

$$3. \mathcal{L}^{-1} \left\{ \frac{1}{s^n} \right\} = \frac{t^{n-1}}{(n-1)!}; \quad n = 1, 2, 3, \dots$$

$$4. \mathcal{L}^{-1} \left\{ \frac{1}{(s - a)^n} \right\} = \frac{e^{at} t^{n-1}}{(n-1)!}$$

$$5. \mathcal{L}^{-1} \left\{ \frac{1}{s^2 + b^2} \right\} = \frac{1}{b} \sin bt$$

$$6. \mathcal{L}^{-1} \left\{ \frac{s}{s^2 + b^2} \right\} = \cos bt$$

$$7. \mathcal{L}^{-1} \left\{ \frac{1}{s^2 - a^2} \right\} = \frac{1}{a} \sinh at$$

$$8. \mathcal{L}^{-1} \left\{ \frac{s}{s^2 - a^2} \right\} = \cosh at$$

$$9. \mathcal{L}^{-1} \left\{ \frac{1}{(s - a)^2 + b^2} \right\} = \frac{1}{b} e^{at} \sin bt$$

$$10. \mathcal{L}^{-1} \left\{ \frac{s - a}{(s - a)^2 + b^2} \right\} = e^{at} \cos bt$$

$$11. \mathcal{L}^{-1} \left\{ \frac{s}{(s^2 + b^2)^2} \right\} = \frac{1}{2b} t \sin bt$$



Centurion  
UNIVERSITY

Shaping Lives...  
Empowering Communities...

# Laplace Transform

## ■ Theorems

<i>Theorem</i>	<i>Description</i>
Definition of Laplace Transform	$\mathcal{L}\{f(t)\} = \int_0^{\infty} e^{-st} f(t) dt = F(s)$
Linear Property	$\mathcal{L}\{af(t) + bg(t)\} = aF(s) + bG(s)$
Derivatives	$\mathcal{L}\{f^{(n)}(t)\} = s^n F(s) - s^{n-1} f(0) - \dots - f^{(n-1)}(0)$
Integrals	$\mathcal{L}\left\{\int_0^{\infty} f(\tau) d\tau\right\} = \frac{1}{s} F(s)$
First Shifting Property	$\mathcal{L}\{e^{at} f(t)\} = F(s - a)$
Second Shifting Property	$\mathcal{L}\{f(t - a)u(t - a)\} = e^{-as} F(s)$



**Centurion**  
**UNIVERSITY**

Shaping Lives...  
Empowering Communities...

## Transform of derivatives

► **Laplace transform of the derivative  $f^n$  of any order :**

Let  $f, f', \dots, f^{(n-1)}$  be continuous for all  $t \geq 0$  and  $f^n$  be piecewise continuous on every finite interval on the semi-axis  $t \geq 0$  then transform of  $f^n$  given by,

$$\mathbf{L}(f^n) = s^n \mathbf{L}(f) - s^{n-1} f(0) - s^{n-2} f'(0) - \dots - f^{(n-1)}(0)$$



**Centurion**  
**UNIVERSITY**

Shaping Lives...  
Empowering Communities...

For first-order derivative:

$$\mathcal{L}\{f'(t)\} = s\mathcal{L}\{f(t)\} - f(0)$$

For second-order derivative:

$$\mathcal{L}\{f''(t)\} = s^2\mathcal{L}\{f(t)\} - sf(0) - f'(0)$$

For third-order derivative:

$$\mathcal{L}\{f'''(t)\} = s^3\mathcal{L}\{f(t)\} - s^2f(0) - sf'(0) - f''(0)$$

For  $n^{\text{th}}$  order derivative:

$$\mathcal{L}\{f^{(n)}(t)\} = s^n\mathcal{L}\{f(t)\} - s^{n-1}f(0) - s^{n-2}f'(0) - \dots - f^{(n-1)}(0)$$

# The Laplace Transform of the **First** Derivative

$$L[f'(t)] = sL[f(t)] - f(0)$$

$$L[f'(t)] = \int_0^{\infty} \underbrace{e^{-st}}_u \underbrace{f(t) dt}_{dv}$$

$$\int u dv = uv - \int v du$$

$$L[f'(t)] = \int u dv = e^{-st} f(t) \Big|_0^{\infty} - \int_0^{\infty} f(t)(-s) e^{-st} dt$$



Centurion  
UNIVERSITY

Shaping Lives...  
Empowering Communities...



**Centurion**  
**UNIVERSITY**

Shaping Lives...  
Empowering Communities...

# The Laplace Transform of the **Third** Derivative

$$L[f'(t)] = sL[f(t)] - f(0)$$

$$L[f''(t)] = s^2 L[f(t)] - sf(0) - f'(0)$$

$$L[f'''(t)] = sL[f''(t)] - f''(0)$$

$$= s \left[ s^2 L[f(t)] - f(0) - f'(0) \right] - f''(0)$$

$$L[f'''(t)] = s^3 L[f(t)] - s^2 f(0) - s f'(0) - f''(0)$$



**Centurion**  
**UNIVERSITY**

*Shaping Lives...  
Empowering Communities...*

# Unit Step Function

$$u(t - a) = 0, t < a$$

$$1, t \geq a$$

$$\begin{aligned} L\{u(t - a)\} &= \int_0^{\infty} e^{-st} u(t - a) dt \\ &= \int_0^a e^{-st} (0) dt + \int_a^{\infty} e^{-st} (1) dt \\ &= \left[ \frac{e^{-st}}{-s} \right]_a^{\infty} = \frac{1}{s} e^{-as} \end{aligned}$$

if  $a = 0$

$$L\{u(t)\} = \frac{1}{s}$$



**Centurion**  
**UNIVERSITY**

Shaping Lives...  
Empowering Communities...

## Application to linear differential equations

We illustrate the Laplace transform technique by a simple example of harmonic oscillator with a time-dependent external force  $f(t) = t$ .

$$\ddot{x} + \omega^2 x = t, \quad (15)$$

$$x(0) = a, \quad \dot{x}(0) = b. \quad (16)$$

The starting point is to Laplace-transform the equation:

$$\int_0^\infty dt e^{-pt} [\ddot{x} + \omega^2 x] = \int_0^\infty dt e^{-pt} t. \quad (17)$$

For the derivatives in the l.h.s. of (17) we use (4), with (16) taken into account. This yields

$$\int_0^\infty dt e^{-pt} [\ddot{x} + \omega^2 x] = (p^2 + \omega^2)X(p) - pa - b, \quad (18)$$

where

$$X(p) = \int_0^\infty dt e^{-pt} x(t). \quad (19)$$

The integral in the r.h.s. of (17) needs to be done explicitly:

$$\int_0^\infty dt e^{-pt} t = -\frac{\partial}{\partial p} \int_0^\infty dt e^{-pt} = -\frac{\partial}{\partial p} \frac{1}{p} = \frac{1}{p^2}. \quad (20)$$

We thus have

$$(p^2 + \omega^2)X(p) - pa - b = \frac{1}{p^2}, \quad (21)$$

and find

$$X(p) = \frac{1 + p^2(pa + b)}{p^2(p^2 + \omega^2)} = \frac{1 + p^2(pa + b)}{p^2(p - i\omega)(p + i\omega)}. \quad (22)$$



Centurion  
UNIVERSITY

Shaping Lives...  
Empowering Communities...

Now we perform the inverse transform

$$x(t) = \frac{1}{2\pi i} \int_C \frac{[1 + p^2(pa + b)] e^{pt}}{p^2(p - i\omega)(p + i\omega)} dp . \quad (23)$$

That is we sum up all the residues of the function

$$\frac{[1 + p^2(pa + b)] e^{pt}}{p^2(p - i\omega)(p + i\omega)} . \quad (24)$$

There are three poles. The two simple poles  $p = \pm i\omega$  come from the l.h.s. of the original differential equation, while the second-order pole at  $p = 0$  is totally due to the particular form of the external force. The residue calculus routine leads to the final answer

$$x(t) = t/\omega^2 + a \cos \omega t + (b/\omega - 1/\omega^3) \sin \omega t . \quad (25)$$



**Centurion**  
**UNIVERSITY**

Shaping Lives...  
Empowering Communities...

**Problem 45.** Use the Laplace transform method to solve for the evolution of a damped harmonic oscillator with a time-independent external force:

$$\ddot{x} + \omega^2 x + \gamma \dot{x} = f_0, \quad (26)$$

$$x(0) = 0, \quad \dot{x}(0) = 0, \quad (27)$$

Find  $x(t)$ ; make sure it is real and does satisfy the equation and the initial conditions. Compare two cases:  $0 < \gamma < 2\omega$  and  $\gamma \geq 2\omega$ .

If we are dealing with a system of linear differential equations, the procedure is the same. We Laplace-transform each of the equations and arrive at a system of *algebraic* equations for the Laplace transforms of the unknown functions. From this system we find each of the Laplace transforms and then perform the inverse transforms.



**Centurion**  
**UNIVERSITY**

Shaping Lives...  
Empowering Communities...

**Problem 46.** Use the Laplace transform method to solve for the evolution of one of two coupled harmonic oscillators (the coupling force is proportional to the relative displacement of the oscillators, the proportionality coefficient  $\lambda > 0$ ):

$$\ddot{x}_1 + \omega_1^2 x_1 = \lambda(x_2 - x_1), \quad (28)$$

$$\ddot{x}_2 + \omega_2^2 x_2 = \lambda(x_1 - x_2). \quad (29)$$

Find  $x_1(t)$  for  $t > 0$  under the following initial conditions:

$$x_1(0) = 0, \quad \dot{x}_1(0) = 0, \quad x_2(0) = 0, \quad \dot{x}_2(0) = 1. \quad (30)$$

Make sure  $x_1(t)$  is real and does satisfy the initial conditions.



**Centurion**  
**UNIVERSITY**

*Shaping Lives...  
Empowering Communities...*

# Laplace Transform


- Applications of the Laplace transform
  - solve differential equations (both ordinary and partial)
  - application to RLC circuit analysis
- Laplace transform converts differential equations in the time domain to algebraic equations in the frequency domain, thus 3 important processes:
  - (1) transformation from the time to frequency domain
  - (2) manipulate the algebraic equations to form a solution
  - (3) inverse transformation from the frequency to time domain



**Centurion**  
**UNIVERSITY**

Shaping Lives...  
Empowering Communities...

## Laplace transform of an impulse 'function' (Dirac delta distribution)



$$L(\delta(t)) = \lim_{\varepsilon \rightarrow 0} \int_0^{\varepsilon} \frac{1}{\varepsilon} e^{-st} dt = \frac{-1}{s\varepsilon} e^{-st} \Big|_0^{\varepsilon} =$$

$$\lim_{\varepsilon \rightarrow 0} \left( \frac{[e^{-s\varepsilon} - 1]}{-s\varepsilon} \right)$$

Writing  $e^x$  as a McLaurin series

$$e^x = 1 + x + \frac{1}{2!}x^2 + \frac{1}{3!}x^3 h.o.t.$$

Writing  $e^{-s\varepsilon}$  as a McLaurin series

$$e^{-s\varepsilon} = 1 + (-s\varepsilon) + \frac{1}{2!}(-s\varepsilon)^2 + \frac{1}{3!}(-s\varepsilon)^3 h.o.t.$$



**Centurion**  
**UNIVERSITY**

Shaping Lives...  
Empowering Communities...

## Laplace transform of impulse (Dirac delta distribution)

$$e^{-s\varepsilon} - 1 = (-s\varepsilon) + \frac{1}{2!}(-s\varepsilon)^2 + \frac{1}{3!}(-s\varepsilon)^3 + h.o.t.$$

$$\frac{e^{-s\varepsilon} - 1}{-s\varepsilon} = 1 + \frac{1}{2!}(-s\varepsilon) + \frac{1}{3!}(-s\varepsilon)^2 + h.o.t.$$

$$L(\delta(t)) = \lim_{\varepsilon \rightarrow 0} \left[ \frac{e^{-s\varepsilon} - 1}{-s\varepsilon} \right] = \lim_{\varepsilon \rightarrow 0} \left[ 1 + \frac{1}{2!}(-s\varepsilon) + \frac{1}{3!}(-s\varepsilon)^2 + h.o.t. \right] = 1$$

$$\boxed{L(\delta(t)) = 1}$$



**Centurion**  
**UNIVERSITY**

*Shaping Lives...*  
*Empowering Communities...*

Solutions to two topics  
are left as an  
assignment.

Submit it before last  
working day of the  
Semester.



**Centurion**  
**UNIVERSITY**

*Shaping Lives...*  
*Empowering Communities...*

**Thank you!**  
**All the best for**  
**Exams !**