Cell Division Mitosis



Reasons for Mitos is There are three main reasons

A. growthB. repair/healingC. as exual reproduction

Cell Division by mitosis

 Some cells divide constantly: cells in the embryo, skin cells, gut lining cells, etc. 7 wee

gut 7 week old embryo







Intestinal Cell

Epithelial Cell

Cell Division

Other cells divide rarely or never.



Brain Cell – Nerve cell

Spinal Cord Cell-Nerve cell





Cardiac Cell (Heart Muscle)

Cell Division

2 kinds of cell division:

I. Mitosis: division of somatic cells

• 2. Meios is : creation of new sex cells





Sperm cells

Human egg cell

Cell Division Vocabulary

 <u>Somatic cell</u> – a body cell; a cell whose genes will not be passed on to future generations.

 <u>Germ cell</u> - a cell that is destined to become a gamete (egg or sperm); a cell whose genes can be passed on to future generations

Cell Division Vocabulary

 <u>diploid (2N)</u> – a cell with 2 chromosome sets in each of its cells; all body (somatic) cells

 <u>haploid (N)</u> – a cell with 1 chromosome set in each of its cells; all gametes (sperm, eggs)

Characteristics of Mitos is

- ADIPLOID CELL WILL GIVE RSE TO ADIPLOID CELL
- GROMOSOME NUMBER FEMANS THE SAME
- THE DNAREMANS IDENTICALLY THE SAME
- ONE CELLOON GIVES RISE TO TWO CELLS (2N)

Cell Cycle

A typical cell goes through a process of growth, development, and reproduction called the cell cycle.

 Most of the cycle is called interphase.



Cell Cycle

 The longest phase in the cell cycle is interphase.

 The 3 stages of interphase are called G₁, S, and G₂.



GAddison Wesley Longman, Inc





Copyright © The Benjamin/Cummings Publishing Company, Inc., from Campbell's BIOLOGY, Third Edition.

Cell Division

- All living cells come from other living cells.
- During mitosis, the nucleus of the cell divides, forming two nuclei with identical genetic information.



Mitos is

Mitos is produces two genetically identical cells. Mitos is referred to in the following stages: prophase, metaphase, anaphase, and telophase.



Prophase

In prophase, the cell begins the process of division.

The chromosomes condense.

(a) DNA in uncondensed form



(b) DNA condensed into duplicated chromosomes





make chromosomes.

chromatin

duplicated chromosom e

cell

Prophase

Nuclear envelope disappears.



Prophase

Centrioles migrate to opposite poles of the cell.

Asters and spindle fibers form.



Aster and the mitotic apparatus in an animal cell

Draw Prophase



Prophase



Metaphase

The chromos omes line up at the equator of the cell (metaphase plate), with the centrioles at opposite ends and the spindle fibers attached to the centromeres.











Draw Metaphase



Anaphase

In anaphase, the centromeres divide.

 At this point, each chromosome goes from having 2 sister chromatids to being 2 separate chromosomes



Anaphase

The spindle fibers contract and the chromosomes are pulled to opposite poles.



Draw Anaphase



Telophase

In telophase the nucleus actually divides. The chromosomes are at the poles of the cell. The nuclear envelope re-forms around the two sets of chromosomes.





Draw Telophase



Cytokines is

The division of the cytoplasm.

 In animal cells, a Cleavage Furrow forms and separates Daughter Cells



Cleavage furrow in a dividing frog cell.

Cytokines is

In plant cells, a Cell Plate forms and separates

Dowahlow Calle



two daughter cells



Cell Plate forming

ANIMAL VS. PLANT MITOSIS

 ANIMAL CELL
Centriole and aster present
Daughter cells separated by cleavage furrow



 PLANT CELL
No visible centriole or aster

> Daughter cells separated by cell plate



















Prophas e





3









Prophase



Metaphas e



3









Prophase



Metaphase



Anaphase









Prophase



Metaphas e



Anaphase



Telophase









Prophase



Metaphas e



Anaphas e



Telophase



C ytokines is



PHASES OF CELLCYCLE-IFMATC



Phases of cell cycle - IPMATC Important **I**mpatient People People <u>M</u>ust May **A**ttack **A**nalyze Tasks Teachers **C**orrectly **C**onstantly