

Water Soluble Vitamins





Vitamin C (Ascorbic acid)

- Water soluble vitamin
- Essential for the formation of collagen which is found in the blood vessels of teeth and bones
- Required for the absorption of iron as it reduces ferric to ferrous form which is easily absorbed

• <u>Functions of Vitamin C</u>

- Essential for the formation of collagen
- Required for wound healing because it helps in the formation of connective tissue
- Essential for the synthesis of norepinephrine a neurotransmitter, activities growth hormone, calcitonin,gastrin releasing peptide
- Excellent antioxidant



Sources & Deficiency

- Food sources-amla ,guava, cashew nut, drumstick leaves,oranges,tomatoes
- Prolonged deficiency causes scurvy in both infants and adults
- Infantile scurvy-loss of apatite , defective growth of bones , defective formation of teeth,swollen gums ,ends of ribs become prominent resulting in scorbutic rosary
- Adult scurvy-fever, susceptibility to infection, delayed wound healing, microcytic hypochromic anaemia develops due to failure of absorption of iron, swollen ulcerated, spongy and bleeding gums, swollen & tender joints, fragile blood vessels, rough and dry skin



Vitamin B1(Thiamine)

- Thiamine is converted to Thiamine pyrophosphate (TPP) which is an coenzyme in carbohydrate metabolism
- TPP is also an essential cofactor for the conversion of tryptophan to niacin
- Involved in the transmission of nerve impulses across the cells
- Sources- whole wheat, millets parboiled rice, bran of cereals, gingelly, groundnut, pork, liver, eggs

Deficiency

• Dry beri-beri-tingling,numbness& burning sensation in hands &legs ,tender calf muscle, complete loss of sensation in hands



Vitamin B1 Deficiency

- Wet beri-beri-enlargement of the heart ,odema in legs and face , palpitations
- Infantile beri-beri-restlessness,enlargement of the heart, breathlessness

• <u>Vitamin B2(Riboflavin)</u>

- Yellow enzyme that is heat stable
- Combined with proteins to form flavo-proteins-FAD,FMN
- Conversion of vitamin B6 and folate to active coenzymes
- Essential for the formation of red blood cells and synthesis of glycogen



Vitamin B2

- Food sources- egg,milk,fish,whole cereals,legumes& greens
- **Deficiency** soreness and burning of the mouth

and tongue, lesiot at the angles of mouth, inflammation of the tongue (glossitis), dry chapped appearance of the lip with ulcers (cheilosis), photophobia, normocytic anaemia, lacrimation

Vitamin B3(Niacin)

- Nictotinamide is essential for the tissue metabolism, the active forms are NAD, NADP
- NADis involved in catabolic reactions &NADP is involved in anabolic



Vitamin B3

- Food sources-dried yeast ,peanuts,whole cereals,legumes,meat, Tryptophan present in dietary proteins is converted to niacin (60mg yields 1 mg)
- **Deficiency** causes a disease Pellagra characteised

by dermatitis, diarrhoea&dementia

- **Dermatitis-** rough skin characterised by reddening,followed by exfoliation leading to parchment
- **Diarrhoea-**structural & absorptive defects in small intestine
- **Dementia** irritability , depression,lossof memory



Vitamin B6(Pyridoxine)

- Functions primarily in protein metabolism
- Three forms that are present in ou body:
- Pyridoxine, Pyridoxal & Pyridoxamine
- Functions:
- Essential for the process of transamination
- Removal of amino group (deamination)
- Removal of carbonyl group (decarboxylation)
- Involved in the conversion of tryptophan to niacin
- Aids in conversion of linoleic to arachidonic acid
- Helps in release of glycogen from liver and muscle



Vitamin B6(Pyridoxine)

- Food sources- meat, pulses, wheat
- Deficiency leads to abnormalities in protein metabolism as poor growth , anaemia, skin leisons decreaseed antibody function, severe deficiency leads to microcytichypochromic anaemia

• <u>VitaminB9 (Folic acid</u>)

- Made of three acids Pteroic,Para amine benzoic acid , glutamic acid
- Functions coenzyme is essential for transferring single carbon & a no.of key compounds are formed: Thymine
- Purines that are essential components of cells
- Conversion of phenylalanine to tyrosine



Vitamin B9

- Sources- Greens, gingelly seeds, cluster beans
- **Deficiency-**bone marrow producing immature cells(megaloblasts) & few matured red bloodcells
- Reduced oxygen carring capacity megaloblastic anaemia, cause neural tube disorders of foetus

<u>VitaminB12(Cyanocobalamin</u>

- Necessary for normal growth and maintenance of healthy nervous tissue
- Involved in DNA synthesis (cell replication)
- Essential for RBC formation
- Facilitates the formation of folate coenzymes needed for nucleic acid synthesis



VitaminB12(Cyanocobalamin

- **Food sources** present in foods only of animal origin,liver sheep,eggs,mutton,milk,and shrimp
- **Deficiency** causes pernicious anaemiacharacterized by very lage immature red blood cells with normal amounts of haemoglobin
- **Requirements-** man/woman-1.0mg/day, children- 0.2-10mg/day