# CARBOHYDRATE METABOLISM



- Introduction
- Classification of Carbohydrates
- Glycolysis
- Glycogenolysis
- Glycogenesis
- Citric acid cycle
- Pentose phosphate pathway
- Applied aspects
- Regulation of Blood glucose



# NUTRITION

Nutrition is defined as "the science of how the body utilizes food to meet requirements for development growth, repair and maintenance"









In plants, Carbondioxide+water

Glucose (stored as starch or converted to cellulose) In Animals, Fat + protein

carbohydrate

# **Biomedical Importance**

Glucose is a major carbohydrate

It is a major fuel of tissues

It is converted into other carbohydrates

Glycogen for storage.

 $\checkmark$ 

 $\checkmark$ 

- Ribose in nucleic acids.
- Galactose in lactose of milk.
- They form glycoproteins & proteoglycans
- They are present in some lipoproteins (LDL).
  - Present in plasma membrane:glycocalyx.
    - Glycophorin is a major intergral membrane glycoprotein of human erythrocytes.

#### Carbohydrates

#### Monosaccharides

#### Disaccharides

Polysaccharide

Oligosaccharides



Depending on carbon atoms •Trioses: Glycerose-Aldo

Dihydroxyacetone-Ketone

•**Tetroses** Erythrose (A)

Erythrulose (K)

•Pentoses: Ribose (A)

Ribulose (K)

•Hexoses: Glucose (A)

Fructose (K)

Depending on aldehyde or ketone group

•Aldoses

•Ketoses

#### •<u>Disaccharides</u> •Maltose

•Sucrose

#### Oligosaccharides •Maltotriose

#### Polysaccharides •Linear - Starch •Branched- Dextrin



The entire spectrum of chemical reactions, occuring in the living system are referred as "Metabolism".

Types of metabolic pathways

- Anabolic pathways: Protein synthesis.
- Catabolic Pathways: Oxidative phosphorylation.
  Amphibolic pathways: Citric acid cycle.

# Food moleculessimpler molecules Amphibolic pathway Anabolic Catabolic P

Proteins, carbohydrates, lipids, nucleic acids etc.

CO2+H2O

Metabolic pathways may be studied at different levels of organisation.

At tissue levelAt subcellular level

# Overview of Carbohydrate

# Metabolism









**Defn**: It is defined as sequence of reactions of glucose to lactate & pyruvate with the production of ATP.

It is derived from greek word *glycose* -sweet or sugar, *lysis*- dissolution.

Site: Cytosolic fraction of cell







#### Glucose-6-phosphate

Phosphoglucose 1

#### Fructose-6-phosphate

Phospho-fructokinase ATP



Fructose-1,6-bisphosphate

#### **STAGE II**





#### 1,3-Bisphosphoglycerate

Phosphoglycerate kinase ADP

#### 3-Phosphoglycerate





#### 2-Phosphoglycerate



#### Phosphoenolpyruvate



#### Pyruvate

# **Bioenergetics in Glycolysis:** Total of 8 ATP is formed in glycolysis. Oxidation of glucose in aerobic condition:38 ATP Anaerobic condition: 2 ATP

### **Biomedical importance of Glycolysis**

- Principal route of metabolism.
- Production of acetyl coA in citric acid cycle.
- Metabolism of fructose & galactose.
- Provides ATP in absence of Oxygen.



- Hemolytic Anaemias: Inherited aldolase A & pyruvate kinase deficiencies.
- Skeletal muscle fatigue
- Inherited Pyruvate dehydrogenase deficiency-Lactic acidosis
- Fast growing cancer cells glycolysis proceeds at faster rate – increased acidic environmentimplication in certain types of cancer.

# **Metabolism of Glycogen**

Major storage form of carbohydrate.

Glycogenesis: occurs in muscle & liver.

### **Biomedical importance**

- Liver glycogen largely concerned with transport & storage of hexose units.
- For maintenance of blood glucose mainly between meals.

#### **<u>Clinical aspects</u>**

#### **<u>Glycogen storage diseases</u>**

Type of disorder Type I (Von Gierke's disease) Type II (Pompe's disease) Type III (Cori's disease)

**Type IV (Andersen's disease)** 

Type V (Mcardle's disease)

**Cause of disorder Glucose-6-phosphatase** deficiency. Acid maltase deficiency. **Debranching enzyme** deficiency. **Branching enzyme** deficiency. **Muscle phosphorylase** deficiency.

**Type VI (Her's disease)** 

Liver phosphorylase deficiency.

**Type VII (Tarui's disease)** 

Phosphofructokinase deficiency

**Type VIII** 

Liver phosphorylase kinase.

•



# **Biomedical importance**

- Final common pathway for oxidation of carbohydrates, lipids, & proteins.
- Major role in gluconeogenesis, transamination, deamination & lipogenesis.
- Vitamins play a key role in this cycle
  Eg; Riboflavin FAD.
  Niacin NAD.
  Thiamine.

Pantothenic acid as a part of co-A. Bioenergetics :12 ATP per cycle.

# **Pentose Phosphate Pathway**

- Alternative route for metabolism of glucose
- It occurs in cytosol
- Sequence of reactions occur in two phases
- 1.Oxidative non reversible phase-Forms NADPH
- 2. Non oxidative reversible phase.- Forms ribose precursors for nucleotide synthesis.

#### Glucose 6-phosphate



#### Ribulose 5-phosphate



# **Biomedical importance**

- Glutathione peroxidase protects erythrocytes against hemolysis.
- Pentose useful in synthesis of DNA & RNA.
- NADPH is required for reductive biosynthesis of fatty acids & steroids.
- NADPH is required in synthesis of amino acids.
- Microsomal cytochrome P450 system brings detoxification of drugs & foreign compounds.

# **<u>Clinical aspects</u>**

Erythrocyte hemolysis

 $\checkmark$ 

 $\checkmark$ 

 $\checkmark$ 

 $\checkmark$ 

 Impairment of generation of NADPH manifests as hemolysis when given drugs like
 Antimalarial- Primaquine aspirin or sulfonamides.

#### (G6 PD) Deficiency:

It makes red cells susceptible to hemolysis X linked inheritance Onset of Anaemia is rapid Mild jaundice

- Defects in Fructose metabolism
- Lack of hepatic fructokinase causes Fructosuria.
- Absence of Hepatic aldolase-Hereditary fructose intolerance.
- Hypoglycemia, vomiting, sweating.
  - Albuminuria, aminoaciduria.

 $\checkmark$ 

Reduced caries incidence.

Fructose & sorbitol in lens associated with diabetic cataract.



- Synthesis of glucose from non carbohydrate compounds is called "gluconeogenesis"
- Site : Mainly occurs in Liver & kidney matrix in cytosol.



indt, c. 2003

# **Regulation of gluconeogenesis**

- Influence of Glucagon.
- Availability of substrates.
- Alcohol inhibits gluconeogenesis.

# Proteoglycans & Glycosaminoglycans

- Seven glycosaminoglycans
- 1 Hyaluronic acid
- 2 Chondriotin sulfate
- 3 Keratan sulfate I
- 4 Keratan sulfate II
- 5 Heparin
- 6 Heparan sulfate
- 7 Dermatan sulfate

### **Mucopolysaccharidoses**

MPS
 MPS I (Hurler syndrome)

MPS II (Hunter syndrome)

MPS IIIA (Sanfilippo A)

MPS IIIB (Sanfilippo B)

MPS IIIC (Sanfilippo C)

Defect Alpha-L-Iduronidase

Iduronate sulfatase

Heparan sulfate N sulfatase

Alpha-Acetylglucosaminidase

Acetyl transferase

MPS IVA (Morquio A)
 Galactose-6-sulfatase

- MPS IVB (Morquio B)
- MPS VI (Maroteaux Lamy syndrome)
- MPS VII (Sly)

 N acetylgalactosamine 4 sulfatase

Beta glucoronidase

Beta galactosidase

### Hunter's syndrome



# **Functions of glycoaminoglycans**

- Structural components of extracellular matrix.
- Act as sieves in extracellular matrix.
- Facilitate cell migration.
- Corneal transparency.
- Anticoagulant (Heparin).
- Components of synaptic & other vesicles.



Oligosaccharide (glycan) covalently attached to their polypeptide backbones. <u>Glycoprotein</u> <u>Functions</u>

Collagen Mucins

Transferrin & Ceruloplasmin Immunoglobulin molecule Alkaline phosphatase Functions Structural molecule Lubricant & protective agent Transport molecule.

Immunity Enzymatic activity

# **Regulation of Blood glucose**

Postabsorptive state: Blood glucose is 4.5-5.5mmol/L.

#### After carbohydrate meal: 6.5-7.2mmol/L

#### During fasting : 3.3-3.9mmol/L

#### **Blood Glucose**



#### Glycogenolysis

#### Gluconeogenesis

<u>Metabolic & hormonal mechanisms</u> <u>regulate blood glucose level</u>

- Maintenance of stable levels of glucose in blood is by
- ✓ Liver.
- Extrahepatic tissues.
- ✓ Hormones .

#### **Liver**

- Freely permeable to glucose via GLUT-2 transporter.
- Passage through cell membrane is rate limiting step.
- Glucose is phosphorylated by hexokinase on entry into cell

#### **Extrahepatic tissues**

- Relatively impermeable to glucose.
- Passage is facilitated through various enzymes.
- It has direct effect on entry of glucose into the cell.



# Role Of Insulin

# **<u>Regulation of blood glucose levels</u>** <u>**Insulin**</u>

Anabolic in response to hyperglycemia

- **Liver** 
  - Stimulates glycogen synthesis, glycolysis, and fatty acid synthesis
- **Muscle** 
  - Stimulates glycogen synthesis

#### Adipose tissue

- Stimulates lipoprotein lipase resulting in uptake of fatty acids from chylomicrons and VLDL
- Stimulates glycolysis for glycerol phosphate synthesis (precursor to triglycerides)

#### Role in insulin in lowering blood glucose





- Produced by A cells of islets of langerhans of pancreas
- Actions opposite to Insulin.
- ✓ Its secretion is stimulated by hypoglycemia.
- It stimulates glycogenolysis & gluconeogenesis from amino acids & lactate.

# **<u>Regulation of blood glucose levels by</u>** <u>Glucagon</u>

Catabolic, in response to hypoglycemia

#### **Liver**

Activates glycogen degradation, gluconeogenesis
 Adipose tissue

Stimulates lipolysis and release of fatty acids





# **Role of thyroid hormone**

It stimulates glycogenolysis & gluconeogenesis. **Hypothyroid** 

- Fasting blood glucose is lowered.
- Patients have decreased ability to utilise glucose.
- Patients are less sensitive to insulin than normal or hyperthyroid patients.

#### Hyperthyroid

- Fasting blood glucose is elevated
- Patients utilise glucose at normal or increased rate



Glucocorticoids are antagonistic to insulin.

Inhibit the utilisation of glucose in extrahepatic tissues.
 Increased gluconeogenesis .



Secreted by adrenal medulla.

✓ It stimulates glycogenolysis in liver & muscle.

 It diminishes the release of insulin from pancreas.

# **Other Hormones**

#### Anterior pituitary hormones

#### **Growth hormone:**

- Elevates blood glucose level & antagonizes action of insulin.
- Growth hormone is stimulated by hypoglycemia (decreases glucose uptake in tissues)
- Chronic administration of growth hormone leads to diabetes due to B cell exhaustion.



Estrogens cause increased liberation of insulin.

Testosterone decrease blood sugar level.

#### **Hyperglycemia**

- Thirst, dry mouth
- Polyuria
- Tiredness, fatigue
- Blurring of vision.
- Nausea, headache,
- Hyperphagia
- Mood change

### **Hypoglycemia**

- Sweating
- Trembling, pounding heart
- Anxiety, hunger
- Confusion, drowsiness
- Speech difficulty
- Incoordination.
- Inability to concentrate



# Glycosuria: occurs when venous blood glucose concentration exceeds 9.5-10.0mmol/L

Fructose-1,6-Biphosphatase deficiency causes lactic acidosis & hypoglycemia..

# **Diabetes Mellitus**

A multi-organ catabolic response caused by insulin insufficiency

**Muscle** 

Protein catabolism for gluconeogenesis

<u>Adipose tissue</u>

Lipolysis for fatty acid release

<u>Liver</u>

- Ketogenesis from fatty acid oxidation
- Gluconeogenesis from amino acids and glycerol <u>Kidney</u>
- Ketonuria and cation excretion
- Renal ammoniagenesis.

# Role of carbohydrates in dental caries

- Fermentable carbohydrates causes loss of caries resistance.
- Caries process is an interplay between oral bacteria, local carbohydrates & tooth surface

Bacteria + Sugars+ Teeth → Organic acids



<u>Role of carbohydrates in periodontal</u>

Abnormal glucose metabolism

disease Excessive carbohydrate intake

**Diabetes Mellitus** 

Obesity

Periodontal disease

Periodontal disease

### **References**

- Text book of Biochemistry –Harper.
  Satyanarayan.
  A C Deb.
- Text book of Physiology –Ganong.
- Text book of Oral Pathology Shafers.
- Principles & practice of Medicine-Davidson.
- Nutrition & oral health The Dental clinics of North America.

Thank you