


STUDENT'S NAME		TOTAL MARKS OBTAINED
CLASS	SUBJECT	
ROLL NO.	DATE	

BUFFER EQUATION / HENDERSON HASELBALCH EQN

This is also referred to as Henderson Haselbalch equation. This equation is used to calculate the pH of the solution or to find out the concentration of H^+ ion.

Consider an acidic buffer containing weak acid, HA and its salt BA.

Dissociation of acid is given by:-



Dissociation of HA is at lesser extent in comparison to its salt. Salt completely dissociates increasing the value of A^- . Using law of Mass Action

$$K_a = \frac{[H^+][A^-]}{[HA]}$$

$[HA]$ - initial concentration

$[H^+]$ - conc of weak acid

$[A^-]$ - " " salt.

Rearranging the equation, we get:-

$$H^+ = \frac{K_a [HA]}{[A^-]}$$

$$H^+ = K_a \times \frac{\text{acid}}{\text{salt}}$$

STUDENT'S NAME	ROLL NO.
SUBJECT	DATE

Taking log

$$-\log [H^+] = -\log K_a - \log \frac{[acid]}{[salt]}$$

Thus $pH = pK_a + \log \frac{[salt]}{[acid]} \rightarrow (\text{Weak Acid})$

pK_a - dissociation exponent (negative logarithm of K_a)

$$pH = pK_a + \log \frac{[salt]}{[base]} \rightarrow (\text{Weak Base})$$

The buffer equation is important in preparation of buffered pharmaceutical solutions. It is satisfactory for calculations with pH range 4 to 10.

$$\frac{[A][H^+]}{[AH]} = K_a$$

$$\frac{[A][H^+]}{[AH]} = K_a$$

$$\frac{[A][H^+]}{[AH]} = K_a$$

BUFFER CAPACITY.

- Buffer Capacity is defined as the resistance offered by the buffer in the change in pH on addition of small amount of acid/base. It is denoted by β .
- Buffer Capacity is also known as Buffer index/ Buffer efficiency / Buffer value.
- Buffer Capacity is also defined as the ratio of small increment caused by strong acids or strong bases causing small change in the pH on its addition.

$$\beta = \frac{\Delta B}{\Delta \text{pH}}$$


ΔB is the small increment in gram equivalents/liter of strong base added to the buffer solution to produce a pH change of ΔpH .

Calculation of Buffer Capacity

1) Consider acetate buffer containing 0.1 mole of acetic acid and 0.1 mole of sodium acetate, in 1 litre of solution.

2) Add 0.01 mole of NaOH.

3) When 0.01 mole of NaOH is added, the concentration of sodium acetate (salt) increases by 0.01 moles/litre, whereas acetic acid (acid) decreases by 0.01 moles/litre.



TOTAL MARKS	STUDENT'S NAME
SUBJECT	CLASS
DATE	ROLL NO.

HAc - Acetic acid. (Acid)

NaAc - Sodium acetate (Salt)

4) This changes in the concentration of salt & acid are represented by buffer equation

$$\text{pH} = \text{pK}_a + \log \frac{[\text{salt}] + [\text{Base}]}{[\text{Acid}] - [\text{Base}]}$$

5) Before adding NaOH, pH of the buffer solution is

$$\text{pH} = 4.76 + \log \frac{0.1 + 0}{0.1 + 0} \\ = 4.76.$$

Thus $\text{pH} = \text{pK}_a$.

6) Buffer capacity decreases rapidly as we add base and is greatest before adding base.
where $[\text{salt}] / [\text{Acid}] = 1$.

7) The more exact equation for calculating Buffer capacity is.

$$\beta = 2.3 C \frac{K_a [H_3O^+]}{(K_a + [H_3O^+])^2}$$

C - total buffer concentration,

K_a - Dissociation constant

$[H_3O^+]$ = hydrogen ion concentration.

* Buffer capacity depends on 2 parameters:-

a) Concentration of salt & acid present

b) Concentration ratio of salt to that of acid

STUDENT'S NAME		TOTAL MARKS OBTAINED
CLASS	SUBJECT	
ROLL NO.	DATE	

BUFFERS IN PHARMACEUTICAL & BIOLOGICAL SYSTEMS

A) Buffers in Pharmaceutical Systems :-

→ Salicylic acid stored in soft glass bottles:-
 Sodium ion leaches out combining with Salicylic acid to form Sodium Salicylate & forming a buffer system. i.e. Salicylic acid & sodium salicylate. This serves as a buffer maintaining the pH & product remains stable.

→ Ephedrine & Ephedrine HCl :-

When HCl is added to Ephedrine fraction of it gets converted to Ephedrine HCl. Mixture HCl & Ephedrine HCl acts as buffer resisting the change in pH of the solution.

→ Buffer system consisting of boric acid, sodium borate and sufficient sodium chloride to make the mixture isotonic & used for ophthalmic solutions in pH range of 7 to 9.

→ Mixture of salts of sodium phosphate with pH 6 to 8. Sodium chloride is added to make it isotonic with body fluids.

B) Buffers in Biological Systems :-

→ In-vivo Biologic Buffer Systems.

a) Blood maintained at pH 7.4 by primary buffers in plasma & secondary

TOTAL MARKS	STUDENT'S NAME
MARKS OBTAINED	SUBJECT
DATE	CLASS

HAc - Acetic acid. (Acid)

NaAc - Sodium acetate (Salt)

4) This changes in the concentration of salt & acid are represented by buffer equation

$$\text{pH} = \text{pKa} + \log \frac{[\text{salt}] + [\text{Base}]}{[\text{Acid}] - [\text{Base}]}$$

5) Before adding NaOH, pH of the buffer solution is

$$\begin{aligned} \text{pH} &= 4.76 + \log \frac{0.1 + 0}{0.1 + 0} \\ &= 4.76 \end{aligned}$$

Thus pH = pKa.

6) Buffer capacity decreases rapidly as we add base and is greatest before adding base.
where $[\text{salt}] / [\text{Acid}] = 1$.

7) The more exact equation for calculating Buffer capacity is:-

$$\beta = 2.3 C \frac{K_a [H_3O^+]}{(K_a + [H_3O^+])^2}$$


C - total buffer concentration,

Ka - Dissociation constant

$[H_3O^+]$ - hydrogen ion concentration.

* Buffer capacity depends on 2 parameters:-

- Concentration of salt & acid present
- Concentration ratio of salt to that of acid



STUDENT'S NAME DRAFT	STUDENT'S NAME
SUBJECT	CLASS
DATE	ROLL NO.

buffers in erythrocytes

- Plasma → Carbonic acid / Bicarbonate & sodium salts of phosphoric acid.
- Erythrocytes → Hemoglobin / Oxyhemoglobin & potassium salts of phosphoric acid.

b) Lacrimal fluid / Tears have pH of about 7.4

→ Urine

- Normal adult urine has pH of about 6. It may be as low as 4.5 or as high as 7.8.
- When pH of urine is below normal values, hydrogen ions are excreted by kidneys and when pH is above 7.4, hydrogen ions are retained by kidneys.