

STUDENT'S NAME	TOTAL MARKS OBTAINED <u>41</u>
CLASS	SUBJECT
ROLL NO.	DATE


## BUFFERED ISOTONIC SOLUTIONS.

- Pharmaceutical solutions that are meant for application to delicate membranes of the body should be adjusted to same osmotic pressure as that of the body fluids.
- An Isotonic solution refers to a solutions having the same osmotic pressure across a semipermeable membrane. eg:- 0.9% NaCl solution
- A Hypertonic solution refers to the one where the concentration of solutes is greater outside the cell than inside it. eg:- 2% NaCl soln.

If the RBCs are suspended in 2% NaCl solution, the water within the cell passes through the cell membrane to dilute the surrounding salt solution. This causes cell shrinkage or crenation.

- A Hypotonic solution refers to the one where the concentration of solute is greater inside the cell than outside it. eg:- 0.9% NaCl soln.

If the RBCs are suspended in 0.9% NaCl solution, the water enters the cell, causing them to swell & finally burst, with the liberation of hemoglobin. This phenomenon is known as hemolysis.



Isotonic      Hypertonic      Hypotonic  
 solute = solute      solute < solute      solute > solute  
 inside      outside      inside      outside      inside      outside

- **Tonicity:** Is the concentration of only the solutes that cannot cross the membrane since these solutes exert osmotic pressure on that membrane.

### \* Measurement of Tonicity

#### 1) Hemolytic method

- The effect of various solution of the drug was observed on the appearance of red blood cells suspended in the solutions.
- method developed by Hunter.
- A quantitative method based on the fact that a hypotonic solution liberates oxyhemoglobin in direct proportion to number of cells hemolyzed.

#### 2) Measurement of slight temperature differences

- It involves measurement of slight temperature differences from differences in the vapour pressure

Freezing point of 0.90% NaCl =  $-0.52^{\circ}\text{C}$ .

STUDENT'S NAME		TOTAL MARKS OBTAINED
CLASS	SUBJECT	
ROLL NO.	DATE	

### 3) Calculating Tonicity using Iso Values.

In case of electrolytes with both the weak & strong types are always greater freezing point depressions are greater than calculated. A new factor,  $L = i K_f$  is introduced.

$$L_{\text{iso}} = \frac{\Delta T_f}{c}$$

$\Delta T_f$  - freezing point depression.  
 $c$  - conc. of solution.


e.g:- Iso value of 0.90% NaCl (0.154M), having  
 $\Delta T_f = 0.52^\circ\text{C}$  is 3.4.

$$L_{\text{iso}} = \frac{0.52}{0.154} = 3.4$$

### \* Methods of Adjusting Tonicity & pH.

There are several methods which are used to render the solution isotonic.

#### 2 Methods



Sodium chloride is added to drug solution to make it isotonic with body fluids.

Water is added to drug solution to make it isotonic with body fluids.

TOTAL MARKS	STUDENT'S NAME
ROLL NO.	CLASS



Class I methods :-

a) Cryoscopic method

b) Sodium chloride equivalent method.

a) Cryoscopic method :-

Freezing point depression of drug solution is determined by the equation

$$\Delta T_f = L_{iso} \times \frac{10}{\text{Mol. wt.}}$$

Example :-

How much NaCl is required to render 100ml of 1% solution of apomorphine HCl isotonic with blood

Given:  $\Delta T_f$  of 0.9% NaCl = 0.52°

$\Delta T_f$  of apomorphine HCl = 0.08°

$\Delta T_f$  of 0.9% NaCl = 0.52°

$\therefore \Delta T_f$  of 1% NaCl = 0.58°

To reduce the freezing point =  $\Delta T_f$  of NaCl -  $\Delta T_f$  ApoHCl  
 $= 0.52 - 0.08$ .

By the method of proportion,

$$\frac{1\%}{x} = \frac{0.58^\circ}{0.44^\circ}$$

$$x = 0.76\%$$

Thus by dissolving 1g of apomorphine HCl & 0.76g of sodium chloride in sufficient  $H_2O$  to make 100 ml solution.

STUDENT'S NAME		TOTAL MARKS OBTAINED
CLASS	SUBJECT	
ROLL NO.	DATE	

## b) Sodium Chloride Equivalent method

Sodium chloride equivalent of drug is the amount of NaCl that has the same osmotic effect of 1g or other weight unit of the drug.

$$\left( E = 17 \times \frac{L_{iso}}{\text{Mol. wt.}} \right)$$

Amount of isotonic agent required to adjust the tonicity.

$$\left( X = \frac{Y}{E} \right)$$

X - grams of isotonic agent required to adjust the tonicity

Y - additional amount of NaCl for isotonicity

E = Gram of NaCl equivalent to 1g of the isotonic agent.

## II) Class II Methods :-

- a) White - Vincent method
- b) The Sprouts method.

### a) White - Vincent method.

- developed equation for calculating the volume V (mls) of osmotic solution prepared by mixing drug with water.

$$(V = w \times E \times 111.1.)$$

w = weight in grams of drug

E = NaCl equivalent.

STUDENT'S NAME	
DEPARTMENT	
ROLL NO.	DATE

STUDENT'S NAME

CLASS

SUBJECT

DATE

## b) The Sprouts Method :-

- Simplification of the White Vincent method.

- Table of values of  $V$  can be constructed by keeping the weight of the drug fixed i.e. 0.3g in 1% solution.

$$\left( \frac{0.3}{1} \times E = V \right)$$

add tannin & borax  $X - X - X$  (bottom of tannin)

$$\left( \frac{X}{E} \right)$$

tannin & borax tannin ratio or excess -  $X$

addition of 1/10 of tannin -  $X$

add 1/10 of borax 1/10 of mordant -  $E$

tannin borax

$\therefore$  about M II and II

bottom tannin - stich (o)

bottom excess -  $E$  (o)

bottom tannin - stich (o)

water add antiseptics not mordants borax bath

1/10 borax mordants tannin -  $E$  (1/10) V

other after wash water

D D

$$(1.11 \times E \times n = V)$$

just a rising of tannin -  $n$

D D  $\therefore$  total drugs  $1.11n = E$