Aquatic Microbiology

Nutrient cycles: Phosphorus, iron and manganese cycles

Phosphorus cycle

- Phosphorus : Mineral nutrient required for plants, animals and microorganisms.
- Microorganisms play a key role:

(i) altering its solubility,

ii) mineralization of organic phosphate into inorganic phosphate,

(iii) oxidation and reduction of phosphorus compounds.

- Uptake of phosphate ions utilized for the synthesis of organic phosphates within the cell, thus a fraction immobilized.
- Upon death of plants , organic phosphate is rapidly released by enzymatic hydrolysis. Phosphate becomes limiting factor.

- The availability of phosphates therefore depends on the degree of solubilization of insoluble phosphates by various organic and inorganic acids produced by microorganisms.
- Soil microorganisms, fungi produce these acids and solubilize insoluble phosphates available to the plants.
- Bacillus, Pseudomonas, Micrococcus, Aspergillus, Penicillium and Fusarium.



Iron cycle

- Iron is an important limiting nutrient for plants, which use it to produce chlorophyll.
- Photosynthesis depends on adequate iron supply. Plants assimilate iron from the soil into their roots.
- Animals consume plants and use the iron to produce hemoglobin, the oxygen transports protein found in red blood cells.
- When animals die, decomposing bacteria return iron to the soil.
- The marine iron cycle is very similar to the terrestrial iron cycle, except that phytoplankton and cyanobacteria assimilate iron.

- Iron fertilization has been studied as a method for sequestering carbon.
- Scientists have hoped that by adding iron to the ocean, plankton might be able to sequester the excess CO2 responsible for climate change.
- However, there is concern about the long term effects of this strategy.
- Iron (Fe) follows a geochemical cycle like many other nutrients.
- Iron is typically released into the soil or into the ocean through the weathering of rocks or through volcanic eruptions.

• The Terrestrial Iron Cycle:

- In terrestrial ecosystems, plants first absorb iron through their roots from the soil. Iron is required to produce chlorophyl, and plants require sufficient iron to perform photosynthesis.
- Animals acquire iron when they consume plants, and iron is utilized by vertebrates in hemoglobin, the oxygen-binding protein found in red blood cells.
- Animals lacking in iron often become anemic and cannot transmit adequate oxygen.
- Bacteria then release iron back into the soil when they decompose animal tissue

- The Marine Iron Cycle: The oceanic iron cycle is similar to the terrestrial iron cycle, except that the primary producers that absorb iron are typically phytoplankton or cyanobacteria.
- Iron is then assimilated by consumers when they eat the bacteria or plankton.
- The role of iron in ocean ecosystems was first discovered when English biologist Joseph Hart noticed "desolate zones," which are regions that lacked plankton but were rich in nutrients.
- He hypothesized that iron was the limiting nutrient in these areas.
- In the past three decades there has been research into using iron fertilization to promote alagal growth in the world's oceans.



Manganese cycle

- Manganese is a chemical element with the symbol Mn and atomic number 25.
- It is not found as a free element in nature; it is often found in minerals in combination with iron.
- In biology, manganese(II) ions function as cofactors for a large variety of enzymes with many functions. Manganese enzymes are particularly essential in detoxification of superoxide free radicals in organisms that must deal with elemental oxygen.

- Manganese also functions in the oxygen-evolving complex of photosynthetic plants.
- While the element is a required trace mineral for all known living organisms, it also acts as a neurotoxin in larger amounts.
- Especially through inhalation, it can cause manganism, a condition in mammals leading to neurological damage that is sometimes irreversible.

Manganese Cycle

