

Taxonomy of Shellfish

Dr. Amrutha Gopan Assistant Professor School of Fisheries Centurion University of Technology and Management Odisha

Introduction

discovers names, determines relationships, classifies and studies the evolution of living organisms.

It is an important branch in biology and is considered to be one of the major subdivisions of biology

Having a broader base than genetics, biochemistry and physiology.

Systematics includes taxonomy

Centurion UNIVERSWO Shaping Lives... Empowering Communities...

term taxonomy is derived from the Greek word 'taxis' - arrangement and 'nomos' - law.

The name taxonomy was first proposed by Candolle (1813).

Taxonomy is defined as the theory and practice of classifying organisms.

In Fisheries science, the taxonomy of aquatic animals could be studied as two major components i.e. **shellfish** taxonomy and **finfish** taxonomy.

The shellfish includes two highly diversified phyla only on Arthropoda and phylum Mollusca. In the provide are named as shellfishes because of the prominent carapace made of chitin as exoskeleton in arthropods and shells made of calcium in molluscs.

These two major phyla are invertebrates.

They show enormous diversity in their morphology, in the habitats they occupy and in their biology.

They include about 6.20 million species with wide habitat preference.

The members of these two phyla have much enturion controlly important groups particularly University Tobsters, shrimps, crabs, oysters, chanks, squids and cuttlefishes.













General characters of the Phylum Arthropoda

NIVERSITY

The Phylum Arthropoda includes spiders, scorpions, ticks, mites, crustaceans, millipedes, centipedes and insects.

This is the most successful phylum of animals both in diversity, distribution and in number of species and individuals.

They show remarkable adaptive features and are successfully adapted to life in water, land and in air.

About 80% of all known animal species are represented in this phylum.

They can survive great extremes of temperature, toxicity, acidity and salinity.

General characters

<u>Centurion</u> <u>Centurion</u> <u>UNIVERSITY</u> <u>Centurion</u> <u>Staping Lives...</u> <u>Empowering Community</u> <u>Some are ectoparasitic and vectors of disease.</u>

- 2. Body have jointed appendages or legs (which are modified to different structures to perform different functions like jaws, gills, walking legs, paddle). There may be 3 pairs, 4 pairs, 5 pairs, many pairs.
- 3. Body is **triploblastic**.

5

- 4. Bilaterally symmetrical.
 - **Organ system** level of organization.
 - Body is divisible into head, thorax and abdomen. NOTE: In some (crustacean and arachnida) body is divisible into cephalothorax (head and thorax is fused) and abdomen.

This is the first group to develop a true head, which contains sense organs and feeding organs specialized for their particular habitats.

ing Live ody is covered with chitinous exoskeleton.

- They are haemocoelomate. Coelom i.e. body cavity is filled with blood or fluid.
- 10. Head bears a pair of compound eyes and antenna.
- 11. Locomotion takes place by jointed appendages.
- 12. Digestive system **complete** with mouth, esophagus, stomach, intestine and rectum with associated glands for digestion.
- 13. Respiration takes place by general body surface or gills (aquatic forms) and trachea or book gills (in terrestrial forms).
- Circulatory system is of open type i.e. do not have blood vessels and enters directly into the body chambers. The blood is colorless.

5. Paired excretory glands (green glands) seen in aquatic crustaceans (shrimps and lobsters) while terrestrial forms whave malphigian tubules as excretory organs.

<u>Chrutien Aquatic forms are ammonotelic, terrestrial forms are universited ic.</u>

- 15. Nervous system is of annelidian type, which consists of brain and ventral nerve cord.
- 16. Unisexual i.e. sexes are separate.

Shaping Lives...

17. Fertilization is internal or external.

18. They are either oviparous or ovoviviparous.

- 19. Development may be **direct or indirect**. During development, larval stages seen in most of the arthropods. The free swimming larva undergoes metamorphosis to become an adult.
- **20. Moulting** or ecdysis, a common feature seen in arthropods. Moulting takes place from very young larval stages onwards. After the moult, the inner body increases in size and is able to adjust its expansion due to the softness of the cuticle.





Among the above four sub-phyla, only the sub-phylum Mandibulata consists of the organisms of our Interests.

