

Centurion university of technology and management

EYE LID TUMOURS

(Basal Cell Carcinoma, Xanthelasma, Hemangioma, Papillomas)



Learning outcome

- Basal cell carcinoma
- Xanthelasma
- Hemangioma
- Papillomas
- Diagnosis, etiology, treatment are included.



Basal cell carcinoma

- Most common malignant tumour of the eyelid.
- Most frequently affects elderly patients.
- Most frequently arises from the lower lid, specially near the inner canthus.
- It does not metastasize.
- Ulcer spreads very slowly, and the growth extends under the skin in all directions and penetrates deeply.





Basal cell carcinoma

CONTINUE...

- UNIVERSITY The pring lives. Empowers Charlen or arises from the basal cell of the epidermis.
- Consists of islands of neoplastic cells (cell nests).
 It is also known as Rodent Ulcer.

Centurion





Rare predisposing conditions

• VINIVERSITY • Vinive Departments who suffer from one of this conditions can develop BCC.

Centurion

(A)Xeroderma pigmentosum(B)Gerlin-Goitz syndrome.



Centurion UNIVERSITY Shaping Lives Drow or ng Connection Exponse of the connection Exponse of the connection

Treatment

Where the tumour is removed surgically.

Radiotherapy.

Cryotherapy, or freezing off the tissue.

Excentration, complete surgical removal of the eyeball and other contents of the eye socket, usually in cases of malignant cancer.

Xanthelesma

Line Vision mon, frequently bilateral condition.

- Found in middle-aged and elderly patients with hyperlipidaemia.
- This is slightly raised, yellow plaque, most commonly found in inner portion of the upper or lower eyelid.
- Most commonly in elderly female.



Centurion UNIVERSITY Ma ping Lives... Empore inter Community et...OV

CONTINUE...

- The ping lives... The ping live
- Signs: yellowish subcutaneous plaque consisting of cholesterol and lipid which located at the medial aspects of eyelids.

Centurion UNIVERSITY Shuping Lives...

Predisposing factors

• Empire in a final commonly, it is spontaneous.

- Associated with diabetes and excess serum cholesterol.
- Histologically, it is the cutaneous deposition of lipid material, being engulfed by histocytes.





Treatment

Destruction by chemicals (trichloroacetic acid), diathermy or photocoagulation.

Recurrence is very common.



Hemangiomas

children.

- It is associated with hemangioma of the choriod and leptomeninges called sturge-weber syndrome.
- It appear bluish when seen through skin, and form swellings which increase in size on crying, on lowering the head.







University Strain Lateral, raised, red lesion which blanches with pressure and may swell on crying.

A large tumour in the upper eyelid may cause mechanical ptosis.





Fig. 1 A capillary hemangioma is an abnormal overgrowth of blood vessels that is sometimes referred to as a "strawberry" birthmark.

Treatment

University

enturio

Large hemangiomas require treatment by intralesional steroid injection, or by superficial radiotherapy.



Papillomas

UNIVERSITY Sh Lyin, Lives E upow pros Communitie ... Own as viral wart.

They are benign growths. Generally it is caused by Human papilloma virus. Signs: A pedunculated or broad based lesion with a characteristic raspberry like surface.



Centurion

CONTINUE...

UNIVERSITY Some Lives of the symptoms: They do not produce any symptoms beyond irritation.

A papilloma is often harmless and does not require treatment.





Treatment

- Excision, surgically removes the papilloma.
- Laser surgery.
- Cryotherapy, or freezing of the tissue.
- Cautery, which involves burning off the tissue.



Books referred

- Clinical ophthalmology, Author: Jack J Kanski ;Publishers: Elsevier ,Edition no:8th edition ,Pg no: 5,7,11,15,16.
- Essentials of ophthalmology.Author: Samar K Bosak.Publishers: Jaypee;Edition no:6th edition,Pg no:135,136,137.
- www.medicalnewstoday.com



Thank you