



ORGANIZATION OF SKELETAL MUSCLE

Suman Kumar Mekap Asst. Professor (Pharmacology) CUTM, Bhubaneswar




ORGANIZATION OF SKELETAL MUSCLE

- All activities that involve movement depend on muscles
- 650 muscles in the human body
- Various purposes formuscles for:
 - Locomotion
 - Upright posture
 - Balancing on two legs
 - Support of internal organs
 - Controlling valves and body openings
 - Production of heat
 - Movement of materials along internal tubes
- Three types of muscles in the human body
 - Skeletal
 - Cardiac
 - Smooth




ORGANIZATION OF SKELETAL MUSCLE

- Skeletal muscles are muscles which are attached to the skeleton.
- •40% of human body mass
- Skeletal muscles are responsible for mainly <u>locomotion</u>, and <u>contraction</u> and *voluntary* <u>relaxation</u>.



ORGANIZATION OF SKELETAL MUSCLE

- Muscle (*whole organ*)
- Fascicle (portion of muscle)
- Muscle Fiber (single muscle cell)
- Myofibril (muscle cell organelle)
- Sarcomere (portion of myofibril)
- Myofilament (part of sarcomere)




- Skeletal muscles are composed of clusters of muscle cells.
 - Muscle fibers
 - Myofibers
 - Myocytes
- A muscle consists of packages of muscle cells called **fascicles**
- A muscle cell is long and spindle shaped





Cell structure

- Muscles cells contain many nuclei
- The plasma membrane \rightarrow *sarcolemma*
- The cytoplasm \rightarrow sarcoplasm
- Length
 - ranges from 0.1cm to more the 30cm in length
- Diameter
 - ranges from 0.001cm to 0.01cm in diameter
- Myofibrils→
 - elongated protein molecules
 - aligned in parallel arrangements
 - extend the full length of the cell.












- The myofibril consists of protein chains called **myofilaments**.
 - Myofilaments have a symmetrical, alternating pattern of thick and thin elements.




Thick myofilament


٠


- consists of a large number of bundled **myosin molecules** aligned in overlapping arrays.
- hexameric proteins with two identical heavy chains and two pairs of different light chains.
 - regulatory light chain (RLC)
 - essential light chain (ELC)






- The thin myofilament (F-actin, filamentous actin)
- made up of two helically intertwined chains of G-actin (globular actin) units.
- Other proteins that bind to the actin molecules:
 - Tropomyosin
 - The Troponin complex \rightarrow made up of three members






• SLIDING FILAMENT MECHANISM :

- The length of skeletal muscle shortens during contraction because the thick and thin filaments slide over one another. The process is known as the sliding filament mechanism.
- The thick filament contains 300 myosin molecules.
- It contain two parts:
- 1. Myosin tail
- 2. Myosin heads
- Myosin tail forms the shaft of the thick filament and heads projects towards the thin filament.
- Thin filament contain actin, troponin and tropomyosin.



- Myosin tail forms the shaft of the thick filament and heads projects towards the thin filament.
- Thin filament contain actin, troponin and tropomyosin.
- At the onset of contraction, the sarcoplasmic reticulum release calcium ions into cytosol
- There they bind to troponin and cause troponintropomyosin complexes to move away from binding site on actin.
- Once the binding sites are free, the repeating sequence of events of the contraction cycle occurs that causes the filaments to slide on each other.



- The contraction cycle consists of 4 steps
- 1. ATP hydrolysis.
- 2. Attachment of myosin to actin to form cross-bridges.
- 3. Power stroke.
- 4. Detachment of myosin from actin.



1. ATP hydrolysis :

- The myosin head includes an ATP- binding site and an ATPase, an enzyme that hydrolyses ATP into ADP and phosphate group.
- •This hydrolysis gives energy to myosin head.
- ADP and a phosphate group remain attached to the myosin head.



2. Attachment of myosin to actin to form cross- bridges:


- The energized myosin head attaches to the myosin binding site on actin and releases the previously hydrolyzed phosphate group.
- When the myosin head attach to actin during contraction, they are referred to as cross-bridges.

Centurion UNIVERSITY Shaping Lives... Empowering Communities...

PHYSIOLOGY OF MUSCLE CONTRACTION

3. Power stroke :

- Once the cross bridges are formed, the power stroke occurs.
- The cross-bridge rotate towards the center of the sarcomere and release the ADP molecule.
- The cross-bridge generates a force which slides the thin filament over the thick filament.



4. Detachment of myosin from actin:

- At the end of power stroke, the cross-bridge remains firmly attached to actin until it binds another molecule of ATP.
- As ATP binds to the ATP binding site on the myosin head, the myosin head detaches from actin.