## Oxidation Reduction Potential

- Oxidation is defined as the process, in which a substance takes up positive, or parts with negative charges, while reduction is the process in which a substance takes up negative or parts with positive charges.
- Oxidation reduction potential (ORP) covers a large pH range. It is an electrochemical test and can easily measured using simple and portable instruments
- $Cu(s) 2e = Cu^{2+}(aq)$  (oxidation, loose electron)
- $Cu^{2+}(aq) + 2e^{-} === Cu(s)$  (Reduction, gain electron)

## **DEFINITION**

- Oxidation Reduction Potential is a measure of the tendency of a chemical species to acquire electrons and thereby be reduced.
- Reduction potential is measured in volts (V), millivolts (mV), or Eh (1 Eh = 1 mV).
- Each species has its own intrinsic reduction potential; the more positive (negative) the potential is , the greater the species' affinity for electrons and tendency to be reduced.

## NERNST EQUATION

For a general reduction reaction,

$$M^{n+}(aq) + ne^- \longrightarrow M_{(s)}$$

Nernst developed an equation, known as: Nernst equation:

$$E = E^{0} - \frac{RT}{nF} \ln \frac{[M_{(s)}]}{[M^{n+(aq)}]}$$

Where,

E is the electrode potential

E° is the standard electrode potential (for 1M solution of metal ions at 298 K)

R is gas constant

T is temperature n is the number of electrons accepted during the change and F is Faraday of electricity (96500 coulombs)

The molar concentration of solids in the pure state is assumed to be unity i.e. [M(s)] = 1. Substituting the values,

R = 8.314 J K-1 mol-1, T = 298 K, F = 96500 coulombs,

the quantity 
$$\frac{2.303RT}{F}$$
 becomes a constant. 
$$\frac{2.303RT}{F} = \frac{2.303 \text{ x } (8.314JK^{-1} \text{ mol}^{-1} \text{ x } (298K)}{(96500 \text{ Cmol}^{-1})} = 0.059$$

Therefore, the Nernst equation for the general reduction reaction at 25°C is:

$$E = E^0 - \frac{0.059}{n} \ln \frac{1}{[M^{n+(aq)}]}$$