

Session_9


Taylor's Series of a Complex Function

If the function $f(z)$ is analytic inside a circle C with centre at a , then for all z inside C ,

$$f(z) = f(a) + f'(a)(z-a) + \frac{f''(a)}{2!}(z-a)^2 + \frac{f'''(a)}{3!}(z-a)^3 + \dots + \frac{f^{(n)}(a)}{n!}(z-a)^n + \dots$$

Proof:

Let z be any point inside the circle C . A circle and radius smaller than that of C such that z is



C_1 is drawn with centre at a an interior point of C_1

Let ' t ' be any point on C_1 , then

$$|z-a| < |t-a| \quad \text{i.e.,} \quad \left| \frac{z-a}{t-a} \right| < 1$$

$$\text{Now, } \frac{1}{t-z} = \frac{1}{(t-a) - (z-a)} = \frac{1}{t-a} \left[1 - \frac{z-a}{t-a} \right]^{-1}$$

Expanding R.H.S. by Binomial theorem as $\left| \frac{z-a}{t-a} \right| < 1$; we obtain

$$\frac{1}{t-z} = \frac{1}{t-a} \left[1 + \left(\frac{z-a}{t-a} \right) + \left(\frac{z-a}{t-a} \right)^2 + \dots + \left(\frac{z-a}{t-a} \right)^n + \dots \right] \quad (2)$$

The series in R.H.S. of (2) converges uniformly since $\left| \frac{z-a}{t-a} \right| < 1$.

Now multiplying both sides of (1) by $\frac{f(t)}{2\pi i}$ and integrating term by term w.r.t t over the circle C_1 , we

obtain

$$\begin{aligned} \frac{1}{2\pi i} \oint_{C_1} \frac{f(t)}{t-z} dt &= \frac{1}{2\pi i} \oint_{C_1} \frac{f(t)}{t-a} dt + \frac{z-a}{2\pi i} \oint_{C_1} \frac{f(t)}{(t-a)^2} dt \\ &+ \frac{(z-a)^2}{2\pi i} \oint_{C_1} \frac{f(t)}{(t-a)^3} dt + \dots + \frac{(z-a)^n}{2\pi i} \oint_{C_1} \frac{f(t)}{(t-a)^{n+1}} dt + \dots \end{aligned} \quad (3)$$

Since $f(t)$ is analytic on and inside C_1 , so by Cauchy integral formula

$$\frac{1}{2\pi i} \oint_{C_1} \frac{f(t)}{t-z} dt = f(z)$$

$$\text{and } f^{(n)}(a) = \frac{n!}{2\pi i} \oint_{C_1} \frac{f(t)}{(t-a)^{n+1}} dt \quad \text{or} \quad \frac{1}{2\pi i} \oint_{C_1} \frac{f(t)}{(t-a)^{n+1}} dt = \frac{f^{(n)}(a)}{n!}$$

∴ From (3) we obtain

$$f(z) = f(a) + (z-a)f'(a) + \frac{(z-a)^2}{2!} f''(a) + \dots + \frac{(z-a)^n}{n!} f^{(n)}(a) \quad (4)$$

which is the required Taylor's series for $f(z)$ about $z = a$.

Maclaurin Series

If $a = 0$, then the Taylor's series (4) is reduced to

$$f(z) = f(0) + zf'(0) + \frac{z^2}{2!} f''(0) + \dots + \frac{z^n}{n!} f^{(n)}(0) + \dots$$

which is known as Maclaurin's series expansion of $f(z)$.