

CONTENTS

- Introduction
- Historical background
- Objectives of breeding
- Flower biology and behavior
- Crossing techniques
- Breeding procedures
- achievements





8 BENEFITS OF TOMATO

Improves vision

Prevents gallstones

Lowers hypertension

Good for digestive health

Helps to manage diabetes

Prevents urinary tract infections

Counters the bad effects of smoking

Reduces cholesterol levels & protects heart

INTRODUCTION



- Common Name : Tomato
- Botanical Name : Solanum lycopersicum formerly Lycopersicon esculentum
- Introduced in India: 19th century by Britishers
- Chromosome No. :2n = 24
- Family: Solanaceae (nightshade)
- Centre of origin :Peru (South America)

OTHER SPECIES

- 1. L. pimpinellifolium Fusarium wilt, early blight resistant.
- 2. L. peruvianum Leaf curl virus resistant.

3. L. cheesmanii - Salt resistant.

4. L. hirsutum - Fruit borer resistant.

5. L. pennellii - Drought tolerant.

AREA, PRODUCTION AND PRODUCTIVITY OF TOMATO

	AREA (000' ha)	PRODUCTION (000'Mt)	PRODUCTIVITY (Mt/ha)
❖ WORLD	4815.71	163029.746	33.9
CHINA	1000	50000	50
❖ INDIA	882.03	18735.9	21.2

Tomato shares 9.4% of total vegetable area and 11.5% of total vegetable production in india.

SOURCE: NHB

HISTORICAL BACKGROUND

- Among the vegetable crops, first commercial F1 hybrid of brinjal was released during 1924 in Japan
- In tomato first F1 hybrid was developed in 1940 in Japan
- At national level first hybrid vigor was reported in chilli during 1933 in by IARI, New Delhi
- First public sector hybrid developed was Pusa Meghdoot in bottle gourd in 1971
- The first hybrid vegetable seeds (Karnataka of tomato and Bharat of bell pepper) were marketed in India by IAHS in 1973

OBJECTIVES OF BREEDING

- > Breeding for earliness.
- > Breeding for increased fruit yield.
- Fruit quality like large round, uniform size, deep red colour and increased shelf life etc.


➤ Breeding for disease resistance like (Fusarium wilt, late blight anthracnose, bacterial wilt).

CONT...

- ➤ Breeding for insect resistance (fruit borer, whitefly etc).
- ➤ Breeding for Abiotic Stresses (cold tolerant, drought tolerant, salt tolerant, low temperature tolerant, herbicide tolerant).
- To breed varieties for prolonged storage and transportation *e.g.*, *flavr Savr*
- > To breed varieties suitable for processing

FLOWER BIOLOGY

The flowers are bisexual, radially symmetric, and consist of 5 parts (sepals, petals, anthers). The calyx is united, at least at the base. The corolla is also united but its shape varies.



FLOWERING BEHAVIOUR

- ANTHESIS: Starts at 6 AM and maximum flower opening till late morning.
- DEHISCENCE: 8 AM 11 AM.
- RECEPTIVITY OF STIGMA: 16 hrs before and 5 days after anthesis.



Plants vary in mating system from completely outcrossing to completely inbreeding (selfing)




Cultivated Tomato

Wild Tomato

- ➤ Cultivated tomatoes are self-pollinating.
- ➤ Pollen shedding often occurs before flowers open.
- ➤ Self-pollinating leads to homozygosis. Inbred lines breed true

CROSSING TECHNIQUES



EMASCULATION PROCEDURE














COLLECTION OF POLLEN







pollination



Hybrid fruits

Breeding procedures

Introduction: Seeds of improved varieties are introduced from one ecological area to another and evaluated. E.g., "Marglobe".

Pure line selection: Arka vikas, Arka saurab.

❖ Back cross method: is commonly utilized in wide crosses or in inter specific gene transfer for resistant to diseases.

Cont.....

- ❖ Pedigree method: has the most common method in tomato. In this method single plant selection is initiated in F2 and is continued through successive generations till pure lines are obtained (up to F6 generations)
- Single seed decent method
- Heterosis breeding
- Mutation breeding

CONT...

Biotechnological methods:

- Breeding for resistance to abiotic stress
- Tomato is sensitive to low and high temp from the stage of germination to ripening. *E.g., Transgenic variety*
- "Flavr Savr" was developed for long shelf life and transportation.

ARKA RAKSHAK





- ✓ First public triple disease resistant tomato F1 hybrid in India.
- ✓ High yielding F1 hybrid giving yield of 90-100 tons per hectare in 140-150 days.
- ✓ Triple disease resistance to tomato leaf curl virus, bacterial wilt and early blight.
- ✓ Suitable for summer, kharif and rabi seasons.

THANK YOU