Chloroquine

Chloroquine is the precedent antimalarial drug. It is used for treating all types of malaria, excluding the one caused by chloroquine-resistant Plasmodium falciparum.

Chemical Structure

Chloroquine

Molecular Formula: C₁₈H₂₆ClN₃

IUPAC Nomenclature: 7-chloro, 4-(4-diethyl amino, 1-methyl) butyl amino quinoline

Chemistry: Chloroquine has quinoline as heterocycle, and chloro group at C7. 4-Dimethyl amino and 1-methyl butyl group substitute the amino group at C4 Therefore, the chemical name of chloroquine is 7-chloro,4-(4-diethylamino,1) methyl butyl amino) quinoline.

Physical Properties: Chloroquine occurs as a white to slightly yellow coloured crystalline powder with no odour and bitter taste. It is soluble in water; and slightly soluble in dilute acids, chloroform, and ether.

1) Crystals of picrate are obtained on treating chloroquine with picric acid

2) A yellow precipitate is obtained when nitric acid solution of chloroquine is warmed with ammonium molybdate solution.

Uses: Chloroquine is used for acute malarial attacks caused by P. vivax, P.malariae, P. ovale, and susceptible strains of P. falciparum. It is also used for suppressive treatment of malaria.

Stability and Storage Conditions: Chloroquine should be stored at room temperature and at 4°C after re-suspension. It loses its colour in sunlight, thus should be stored in tightly-closed light-resistant containers.

Types of Pharmaceutical Formulations: Syrups, tablet, and injection.

Brand Names: Amoquin, Antima, Bitaquine, Clokit, and Ciplaquin.

Amodiaquine 9.1.3.2.

a 4-aminoquinoquinoline derivative containing Amodiaguine is inflammatory properties.

Molecular Formula: C20H22ClN3O

IUPAC Nomenclature: 4-[(7-chloroquinolin-4-yl) amino]-2-(diethylaminomethyl)phenol

Properties: Amodiaquine occurs as a yellow coloured crystalline powder with no odour and bitter taste. It is soluble in water; and very slightly soluble in ethanol.

Uses: Amodiaquine is an antimalarial drug used for treating acute malarial attacks having symptoms like fever, chills, and sweating.

Storage Conditions: Amodiaquine should be stored in air-tight containers, away from children.

Types of Pharmaceutical Formulations: Tablet and suspension.

Brand Names: Basoquine, Camoquin, Eriquin, and Flavoquin.

9.1.3.3. Primaquine

Primaquine is an aminoquinoline which is indicated orally for radically curing and preventing relapse of vivax and ovale malaria after treatment with blood schizontocide.

Minima Salemapici 9)

Molecular Formula: C15H21N3O

IUPAC Nomenclature: 4-N-(6-methoxyquinolin-8-yl)pentane-1,4-diamine

Properties: Primaquine occurs as an orange-red coloured crystalline powder with a bitter taste and no odour. It is soluble in water; and insoluble in alcohol, chloroform, and ether.

Uses: Primaquine is used for treating malaria caused by P. ovale and P. vivax.

Stability and Storage Conditions: Primaquine should be stored in tightly-closed containers, protected from light and moisture.

Types of Pharmaceutical Formulation: Tablet.

Brand Names: Malirid and Aralon.

9.1.3.4. Proguanil

Proguanil is a biguanide compound which forms cycloguanil (an anti-malarial) by getting metabolised in the body. It is generally indicated with atovaquone.

Molecular Formula: C₁₁H₁₆ClN₅

IUPAC Nomenclature: (1E)-1-[amino-(4-chloroanilino)methylidene]-2-propan-2-ylguanidine

Chemistry: Proguanil is a biguanidine derivative with its N^1 -nitrogen getting substituted by p-chlorophenyl group and N^5 -nitrogen getting substituted by isopropyl group. it converts into cycloguanil (an active principle exhibiting antimalarial activity) within the liver.

Physical Properties: Proguanil occurs as a white crystalline powder with a bitter taste and no odour. It is slightly soluble in water; and insoluble in chloroform.

Chemical Properties: A white precipitate is obtained on treating aqueous solution of proguanil with potassium ferrocyanide.

Uses: Proguanil is used for preventing and suppressing malaria caused by various strains of *P. falciparum* and other species of *Plasmodium*,

Stability and Storage Conditions: Proguanil should be stored at 25°C (77°F). Temperature changes are permitted between 15-30°C. It gets affected by light, thus it should be stored in tightly-closed light-resistant containers.

Types of Pharmaceutical Formulations: Tablet and syrup.

Brand Names: Malarone, Laveran, and Progunal.