3.1.3.1. Sulphadiazine

Sulfadiazine is a short-acting bacteriostatic and a synthetic pyrimidinyl sulfonamide derivative.

Molecular Formula: C10H10N4O2S

IUPAC Nomenclature: 4-amino-N-pyrimidin-2-ylbenzenesulfonamide

Physical Properties: Sulfadiazine occurs as a white to slightly yellow coloured crystalline powder. It is soluble in boiling water; faintly soluble in ethanol and acetone; and insoluble in water, chloroform, and ether. It solubilises easily in dilute hydrochloric acid, sodium hydroxide solution, and ammonia solution.

Chemical Properties:

1) Diazotisation Reaction: Sulphadiazine undergoes diazotisation and gives a diazo compound on being treated with sodium nitrite in the presence of concentrated hydrochloric acid at 0-10°C.

p-(pyrimid-2-yl amino sulphonyl) benzene diazonium chloride

 Sulphadiazine can be identified by the presence of a purple brown coloured precipitate on being dissolved in 0.1N sodium hydroxide and 1% copper sulphate solution.

Uses: Sulphadiazine can be used for the treatment of upper respiratory tractinfections, otitis media, *Meningococcal* meningitis, boils carbuncle, puerperafever, urinary tract infections, acute dysentery, etc. It is also used for treating

infections caused by Haemolytic streptococcus, Meningococcis, Pneumococcus, Neisseria gonorrhoea, and E. coli.

Stability and Storage Conditions: Sulphadiazine on exposure to light becomes dark, thus should be stored away from light and moisture in well-closed containers at room temperatures ranging between 15-30°C.

Types of Pharmaceutical Formulations: Tablets and creams.

Brand Names: Atrina, Cotrizine, and Trimozin.

3.1.3.2. Sulphaguanidine

Sulphaguanidine (a guanidine derivative of sulphanilamide) is a veterinary medicine. It is suitable for treating dysentery and other enteric infections.

Molecular Formula: C7H10N4O2S

TUPAC Nomenclature: 2-(4-aminophenyl) sulfonylguanidine

Properties: Sulphaguanidine occurs as a white, fluffy powder. It is practically insoluble in water.

Chemical Properties: Sulphaguanidine undergoes diazotisation reaction being treated with sodium nitrite in the presence of concentrated hydrochloric acid at 10°C temperature.

In this reaction, the aromatic primary amino group converts into diazo group.

p-(guanidinyl sulphonyl) benzene diazonium chloride

Uses: Sulphaguanidine is used as a raw material of sulfadiazine drugs, as auxiliary material for other drugs, and for treating bacillary dysentery.

Storage Conditions: Sulphaguanidine should be stored in tightly-closed containers in a dry and well-ventilated place.

Types of Pharmaceutical Formulation: Tablet.

Brand Names: Ganiden, Guanimycin, and Entrogen.

Phthalylsulphathiazole 3.1.3.3.

Phthalylsulphathiazole is a broad spectrum antimicrobial agent used for treating various infections including intestinal.

Molecular Formula: C₁₇H₁₃N₃O₅S₂

Nomenclature: **IUPAC**

2-[[4-(1,3-thiazol-2,

ylsulfamoyl)phenyl]carbamoyl]benzoic acid

Properties: Phthalylsulphathiazole occurs as a colourless, amorphous powder having a characteristic odour.

Uses: Phthalylsulphathiazole is prescribed for treating dysentery, gastrointestinal infection, colitis, and intestinal surgery.

Storage Conditions: Phthalylsulphathiazole should be stored in a cool, moisturefree place, out of the reach of children.

Types of Pharmaceutical Formulation: Tablets.

Brand Names: Diarrocalmol, Ftalasol, and Liracol.

Succinylsulphathiazole

N-succinylsulphathiazole is a member of 1, 3-thiazoles and a derivative of sulphathiazole.

Molecular Formula: C₁₃H₁₃N₃O₅S₂

IUPAC Nomenclature: 4-oxo-4-[4-(1,3-thiazol-2-ylsulfamoyl)anilino]butanoic acid

Properties: Succinylsulphathiazole occurs as a white or yellow coloured crystalline powder. It is soluble in alkali hydroxide. It solubilises in sodium bicarbonate solutions with carbon dioxide evolution. It is sparingly soluble in alcohol and acetone; and insoluble in chloroform and ether.

Uses: Succinylsulphathiazole shows antibacterial activity in GIT.

Storage Conditions: Succinylsulphathiazole should be stored at room temperature.

Types of Pharmaceutical Formulation: Tablets.

Brand Names: Guanicar, Cremosuxidine, and Kaoxidine.