Half Range Expansion of a Fourier series:-

Even and odd functions:-

Even Function:-

A function f(x) is called even if f(-x) = f(x) for all value of x.

Properties of even functions:-

- The graph of an even function is always symmetrized about the y-axis.
- f(x) contains only even power of x and may contains only cosx, sec x.
- Sum of two even function is even.
- Product of two even function is even.
- $\int_{-a}^{a} f(x) dx = 2 \int_{0}^{a} f(x) dx$

Odd Function:-

A function f(x) is called odd if f(-x) = -f(x), $\forall x \in \mathbb{R}$

- The graph of an odd function is always symmetric about the origin and lie in opposite quadrant (1st and 3rd).
- f(x) contains only odd power of x and may contains only sin x, cosec x.
- $\int_{-a}^{a} f(x) dx = 0.$
- Sum of two odd function is odd function .
- Product of an odd function and even function is an odd function.
- Product of two odd function is an even function .

Neither Even nor Odd functions:-

A function f(x) which is not satisfying the even and odd condition is called neither even nor odd function.

Some Useful Formulas:-

• Sin $n\pi = \cos(n+1/2) \pi = 0 \forall n \in \mathbb{Z}$

i.e. $\sin \pi = \sin 2\pi = \sin (3\pi = \sin (-\pi) = \sin (-2\pi) = \dots = 0$

i.e. $\cos \pi/2 = \cos 3\pi/2 = \cos 5\pi/2 = \dots = 0$

• $\cos n\pi = \sin(n+1/2) \pi = (-1)^n \forall n \in \mathbb{Z}$

Cos $\pi = \cos 3\pi = \cos 5\pi = = -1$ Sin $\pi/2 = \sin 5\pi/2 = \sin 9\pi/2 = ... = 1$ Cos $0 = \cos 2\pi = \cos 4\pi = --- = 1$ Sin $3\pi/2 = \sin 7\pi/2 = \sin 11\pi/2 = ... - 1$

• If x=x₀ is the part of the finite discontinuity then sum of the fourier series

$$f(x) = \frac{1}{2} \left[\lim_{h \to 0} f(x_0 - h) + \lim_{h \to 0} f(x_0 + h) \right]$$

$$f(x) = \frac{1}{2} \left[f(x_0 - 0) + f(x_0 + 0) \right]$$

Even and odd Extensions of a Function:-

Let f be a function given on the interval (0,a),we define the even/odd extension of f to be even/odd functions on the interval (-a,a),which coincides with f on the half interval (0,a).

Even Extension:-

$$f_E = even f = \begin{cases} f(-x), & -a < x < 0 \\ f(x), & 0 < x < a \end{cases}$$

Odd Extension:-

 $f_0 = odd f =$

$$\begin{cases} -f(-x), -a < x < 0 \\ f(x), \quad 0 < x < a \end{cases}$$

Half Range Expansion of a Fourier series:-

Suppose a function is defined in the range(0,L), instead of the full range (-L,L).Then the expansion f(x) contains in a series of sine or cosine terms only .The series is termed as half range sine series or half range cosine series.

 If f(x) is taken to be an odd function, its Fourier series expansion will consists of only sine terms. Hence the Fourier series expansion of f(x) represents "Half range expansion of Fourier sine series".

i.e f(x) =
$$\sum_{n=1}^{\infty} b_n \sin \frac{n\pi x}{L}$$
, (0b_n = \frac{2}{L} \int_0^L f(x) \sin \left(\frac{n\pi x}{L}\right) dx.

• If f(x) is taken to be an even function , then its Fourier series expansion will consists of constant and cosine terms. Therefore the Fourier series expansion of f(x) represents "Half range expansion of Fourier series" i.e.

$$f(\mathbf{x}) = f(\mathbf{x}) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left(a_n \cos \frac{n\pi x}{L} \right)$$

where $a_0 = \frac{2}{L} \int_0^L f(x) dx$

$$a_n = \frac{2}{L} \int_0^L f(x) \cos \frac{n\pi x}{L} dx$$
, for all $(n \ge 1)$ $n \in \mathbb{Z}$

Here

 $a_0 \ a_n \ b_n$ are called coefficients half range Fourier cosine series.

Example – **3**:-Determine the half range Fourier series sine series for

$$f(x) = e^x \text{ in } 0 < x < 1$$

Solution:-

Given that $f(x) = e^x 0 < x < 1$ we know that half rage Fourier sine series is

$$f(x) = \sum_{n=1}^{\infty} \left(b_n \sin \frac{n\pi x}{L} \right)$$
 here $l = 1$

Now $b_n = 2/L \int_0^L f(x) \sin \frac{n\pi x}{L} dx$

$$=2\int_{0}^{1} e^{x} \sin n\pi x \, dx$$
$$=2\left[\frac{e^{x}}{1+n^{2}\pi^{2}}(\sin n\pi x - n\pi \cos n\pi x)\right]_{0}^{1} = \left[\frac{2n\pi}{1+n^{2}\pi^{2}}(-n\pi e(-1)^{n} + n\pi)\right]$$

Therefore $f(x) = 2\pi \sum_{n=1}^{\infty} \frac{n(1-e(-1)^n)}{1+n^2\pi^2} \sin n\pi x$

Example – **4**:-Determine the half range Fourier cosine series $f(x)=\sin x$ in $0 < x < \pi$ **Ans**:-Given that $f(x) = \sin x$ is defined in $(0, \pi)$ ie $L = \pi$

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left(a_n \cos \frac{n\pi x}{L} \right)$$
$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} (a_n \cos nx) \quad after putting \ L = \pi$$

Now

$$a_{0} = \frac{2}{\pi} \int_{0}^{\pi} f(x) dx = \frac{2}{\pi} \int_{0}^{\pi} \sin x \, dx = \frac{2}{\pi} [-\cos x]_{0}^{\pi} = \frac{2}{\pi}$$
$$a_{n} = \frac{2}{\pi} \int_{0}^{\pi} f(x) \cos nx \, dx = \frac{2}{\pi} \int_{0}^{\pi} \sin x \cos nx \, dx$$
$$= -\frac{2}{\pi (n^{2} - 1)} [-(-1)^{n} - 1] \quad provided \ n \neq 1$$

By definition
$$a_n = \begin{cases} 0 \text{ if } n \text{ is odd} \\ -\frac{4}{\pi(n^2-1)} \text{ if } n \text{ is even} \end{cases}$$

Hence, the half range cosine series is given by

$$f(x) = \frac{2}{\pi} - \frac{2}{\pi} \sum_{n=2}^{\infty} \frac{\cos nx}{n^2 - 1} (1 + (-1)^n)$$

i.e $\sin x = \frac{2}{\pi} - \frac{4}{\pi} (\cos \frac{2x}{1*3} + \cos \frac{4x}{3*5} + \cos \frac{6x}{5*7} + \dots)$ is the cosine series.

required cosine series.

Exercise 2.3

Are the following functions odd, even, neither odd nor even

(a) $x^{2} \cos x$ (b) $\sin x + \cos x$ (c) $f(x) = e^{x^{2}}$ (d) $\ln x$ (e) $e^{|x|}$ (f) $f(x) = x^{4}$ ($0 < x < 2\pi$)

(g)
$$f(x) = \begin{cases} \cos^2 x , if -\pi < x < 0 \\ \sin^2 x, if \ 0 < x < \pi \end{cases}$$
 (h) $e^{-|x|}$ (- $\pi < x < \pi$)

2. Determine half range Fourier sine series for $f(x)=x(\pi-x)$ in $0 < x < \pi$

hence deduce $\frac{1}{1^3} + \frac{1}{3^3} + \frac{1}{5^3} + \dots = \pi^3/3^2$.

3. Determine half range Fourier sine series for $f(x)=e^x$ in $0 < x < \pi$

4.Determine the half range Fourier sine series for

$$f(x) = \begin{cases} x , & 0 < x < \frac{\pi}{2} \\ \pi - x , & \frac{\pi}{2} < x < \pi \end{cases}$$

5. Determine the half range Fourier cosine series for

$$f(x) = \begin{cases} x , & 0 < x < \frac{\pi}{2} \\ \pi - x , \frac{\pi}{2} < x < \pi \end{cases}$$

6.Find half-range Fourier cosine series for f(x)=x in $(0 < x < 2\pi)$