

HEPATITIS A,B,C,D,E,F



Centurion
UNIVERSITY

Shaping Lives...
Empowering Communities...

By
Ms. RUPALI RUPASMITA ROUT
ASSISTANT PROFESSOR

INTRODUCTION:

Hepatitis refers to inflammation of the liver, which can be caused by a variety of factors, including viral infections, alcohol consumption, certain medications, and autoimmune disorders.

TYPES: Hepatitis A, B, C , D, E, F.

ETIOLOGY:

Hepatitis A: This type of hepatitis is caused by the hepatitis A virus (HAV) and is typically spread through contaminated food or water. Poor hygiene and inadequate sanitation are often the primary factors that contribute to the spread of this virus.

Hepatitis B: This type of hepatitis is caused by the hepatitis B virus (HBV) and is primarily spread through contact with infected bodily fluids, such as blood or semen. This can occur through unprotected sex, sharing needles or other drug injection equipment, or from mother to baby during childbirth.

Hepatitis C: This type of hepatitis is caused by the hepatitis C virus (HCV) and is also primarily spread through contact with infected bodily fluids, such as blood or semen. Sharing needles or other drug injection equipment is a common way that this virus is spread.

Hepatitis D: This type of hepatitis is caused by the hepatitis D virus (HDV) and can only occur in people who are already infected with hepatitis B. The virus is primarily spread through contact with infected blood or bodily fluids.

Hepatitis E: This type of hepatitis is caused by the hepatitis E virus (HEV) and is primarily spread through contaminated food or water. This type of hepatitis is more common in developing countries with poor sanitation.

Hepatitis F: Cause of hepatitis F is unknown. and is primarily spread through contaminated food or water. This type of hepatitis is more common in developing countries with poor sanitation.

OTHER CAUSES:

Alcohol abuse: Excessive alcohol consumption can cause inflammation of the liver and lead to hepatitis.

Autoimmune disorders: The body's immune system can attack the liver, leading to inflammation and hepatitis.

Toxins: Exposure to certain chemicals and toxins, such as drugs, alcohol, and industrial chemicals, can also cause hepatitis.

Other infections: Hepatitis can also develop as a result of other infections, such as cytomegalovirus, Epstein-Barr virus, and yellow fever.

PATHOPHYSIOLOGY:

Virus entry: The hepatitis virus enters the body through various routes, depending on the specific virus. For example, hepatitis A, E and F viruses are usually contracted by consuming contaminated food or water, while hepatitis B, C, and D viruses are transmitted through blood or other bodily fluids.

Targeting liver cells: Once the virus enters the body, it travels to the liver, which is the primary target organ. The virus then targets liver cells, known as hepatocytes, and begins to replicate within them.

Immune response: As the virus replicates in the liver cells, the body's immune system detects the presence of the virus and shows an inflammatory response. The immune system sends white blood cells to the liver to attack the infected cells, which can cause further damage to the liver tissue.

Inflammation: The immune response triggers inflammation in the liver, which can lead to the development of hepatitis. Chronic inflammation can cause ongoing liver damage, and over time, this damage can lead to cirrhosis or liver cancer.

CLINICAL MANIFESTATION:

- Fatigue
- Abdominal pain
- Nausea
- Vomiting
- Jaundice (yellowing of the skin and eyes)
- High fever.

DIGNOSIS:

Blood tests: Blood tests can be used to detect the presence of the hepatitis virus and to determine the extent of liver damage. The tests may include:

1. **Hepatitis serology:** This blood test detects antibodies to the hepatitis virus, which can indicate whether a person has been infected with the virus.
 2. **Liver function tests (LFTs):** These blood tests measure the levels of certain enzymes and proteins in the blood that are produced by the liver. Abnormal levels can indicate liver damage.
 3. **Viral load tests:** These blood tests measure the amount of virus present in the blood. This can help doctors determine the severity of the infection and monitor the effectiveness of treatment.
- **Imaging tests:** Imaging tests, such as ultrasound, CT scans, or MRI scans, can be used to examine the liver and detect any abnormalities, such as scarring or liver tumors.
 - **Liver biopsy:** A liver biopsy involves taking a small sample of liver tissue for examination under a microscope. This can help doctors determine the extent of liver damage and rule out other liver diseases.

- **Other tests:** In some cases, other tests may be done to diagnose hepatitis, such as urine tests, stool tests, or tests to check for other infections that may be contributing to the liver damage.

PHARMACOLOGICAL MANAGEMENT:

- **Hepatitis A:** There is no specific treatment for hepatitis A, but most people recover on their own within a few weeks. Rest, adequate hydration, and avoiding alcohol and certain medications that can be harmful to the liver are recommended.
- **Hepatitis B:** Antiviral medications are used to treat chronic hepatitis B. These drugs work by reducing the amount of virus in the body, which can slow or stop the progression of liver damage. Examples of antiviral medications used to treat hepatitis B include tenofovir, entecavir, and lamivudine.
- **Hepatitis C:** Antiviral medications are also used to treat hepatitis C. These drugs work by blocking the replication of the virus, which can cure the infection in many cases. Examples of antiviral medications used to treat hepatitis C include sofosbuvir, ledipasvir, and daclatasvir.
- **Hepatitis D:** There is no specific treatment for hepatitis D, but antiviral medications may be used to treat concurrent hepatitis B infections.
- **Hepatitis E:** There is no specific treatment for hepatitis E, but most people recover on their own within a few weeks. Rest, adequate hydration, and avoiding alcohol and certain medications that can be harmful to the liver are recommended.

NON- PHARMACOLOGICAL MANAGEMENT:

- **Get vaccinated:** Vaccines are available for hepatitis A and B, and getting vaccinated can provide protection against these viruses.
- **Practice good hygiene:** Wash your hands thoroughly with soap and water after using the bathroom, changing diapers, or handling food.
- **Avoid contaminated food and water:** Be cautious when traveling to areas with poor sanitation, and avoid consuming raw or undercooked shellfish, meat, and fish.
- **Practice safe sex:** Use condoms to reduce the risk of transmitting hepatitis B and C during sexual activity.
- **Avoid sharing personal items:** Do not share needles, razors, toothbrushes, or other personal items that could be contaminated with blood or bodily fluids.
- **Be cautious with tattoos and piercings:** Ensure that any equipment used for tattoos or piercings is properly sterilized to avoid exposure to hepatitis B and C.
- **Manage underlying medical conditions:** If you have an underlying medical condition that can increase your risk of hepatitis, such as HIV, seek appropriate medical care and follow your healthcare provider's recommendations for preventing hepatitis.