

MEGALOBLASTIC ANEMIA



Centurion
UNIVERSITY

Shaping Lives...
Empowering Communities...

By
Ms. RUPALI RUPASMITA ROUT
ASSISTANT PROFESSOR

INTRODUCTION:

Megaloblastic anemia is a type of anemia characterized by the presence of abnormally large red blood cells, known as megaloblasts, in the red bone marrow. It is primarily caused by a deficiency of vitamin B12 (cobalamin) or folate (vitamin B9), which are essential for the production of healthy red blood cells.

ETIOLOGY/CAUSES:

- **The most common causes of megaloblastic anemia include:**
- **Vitamin B12 deficiency:** This can occur due to inadequate dietary intake of vitamin B₁₂, malabsorption of the vitamin (often associated with conditions like pernicious anemia or gastrointestinal disorders), or impaired utilization of vitamin B₁₂ by the body.
- **Folate deficiency:** Inadequate dietary intake of folate, poor absorption in the intestines, increased requirements (e.g., during pregnancy), or certain medications (such as methotrexate) can lead to folate deficiency.
- **Causes of Vitamin B12 Deficiency:**
- **Pernicious Anemia:** This is an autoimmune condition in which the body's immune system attacks the cells in the stomach that produce intrinsic factor, a protein required for vitamin B12 absorption.
- **Dietary Insufficiency:** Strict vegetarians and vegans who do not consume animal products, which are the main sources of vitamin B12, may develop a deficiency if they do not supplement their diet adequately.
- **Malabsorption:** Certain gastrointestinal disorders, such as celiac disease, Crohn's disease, and gastric bypass surgery, can impair the absorption of vitamin B12 from the diet.

- **Parasitic Infections:** Infections caused by parasites like fish tapeworms (*Diphyllobothrium latum*) can deplete vitamin B12 levels by consuming it from the intestines.
- **Medications:** Long-term use of certain medications, such as proton pump inhibitors (PPIs) and metformin, can interfere with the absorption or utilization of vitamin B12.
- **Causes of Folate Deficiency:**
- **Poor Dietary Intake:** Inadequate consumption of foods rich in folate, such as leafy green vegetables, citrus fruits, and legumes, can lead to folate deficiency.
- **Malabsorption:** Conditions affecting the intestines, such as celiac disease, inflammatory bowel disease, or surgical removal of a significant portion of the small intestine, can impair folate absorption.
- **Alcoholism:** Excessive alcohol consumption can interfere with folate absorption, utilization, and increase folate excretion.
- **Pregnancy and Lactation:** Folate requirements are increased during pregnancy and breastfeeding, and inadequate intake can lead to deficiency.
- **Medications:** Certain medications, such as methotrexate (used to treat cancer and autoimmune diseases), can interfere with folate metabolism.

PATHOPHYSIOLOGY:

- **Red Blood Cell Formation:** The bone marrow continuously produces new red blood cells through a process called erythropoiesis. This process involves the division and maturation of precursor cells known as erythroblasts.

- **DNA Synthesis:** During erythropoiesis, DNA synthesis is crucial for the proper division and maturation of erythroblasts. DNA provides instructions for cell growth, replication, and function.
- **Vitamin B12 and Folate Role:** Vitamin B12 and folate are essential for the production of nucleotides, which are the building blocks of DNA. Vitamin B12 is necessary for converting folate into its active form, methylfolate, which participates in DNA synthesis.
- **Vitamin B12 Deficiency:** When there is a deficiency of vitamin B12, such as in pernicious anemia or inadequate dietary intake, the conversion of folate to its active form is impaired. As a result, there is a decrease in the availability of active folate for DNA synthesis.
- **Folate Deficiency:** In cases of folate deficiency, which can be caused by poor dietary intake or malabsorption, there is a direct lack of folate available for DNA synthesis.
- **Impaired DNA Synthesis:** Without sufficient vitamin B12 or folate, the DNA synthesis process becomes compromised. DNA replication and cell division are disrupted, resulting in the production of larger and structurally abnormal erythroblasts called megaloblasts.
- **Large Red Blood Cells:** These megaloblasts mature into abnormally large red blood cells known as macrocytes. These macrocytes are ineffective in carrying oxygen and have a shorter lifespan, leading to anemia.
- **Anemia Symptoms:** The reduced number of functional red blood cells and the abnormal shape of macrocytes result in the characteristic symptoms of

megaloblastic anemia, including fatigue, weakness, pale skin, shortness of breath, and other related symptoms.

CLINICAL MANIFESTATION:

- **Fatigue and Weakness:** Feeling tired and lacking energy is a common symptom of anemia in general, including megaloblastic anemia. The body's reduced ability to deliver oxygen to tissues and organs can result in fatigue and weakness.
- **Pale Skin:** Anemia often leads to paleness or a noticeably lighter complexion. This occurs because the reduced number of red blood cells affects the redness of the skin.
- **Shortness of Breath:** Insufficient red blood cells can limit the amount of oxygen transported to the lungs and other organs. This can cause difficulty in breathing, especially during physical exertion.
- **Rapid Heartbeat (Tachycardia):** The heart may beat faster in an attempt to compensate for the reduced oxygen-carrying capacity of the blood.
- **Dizziness and Lightheadedness:** Inadequate oxygen supply to the brain can result in feelings of dizziness, lightheadedness, or even fainting spells.
- **Glossitis:** Megaloblastic anemia can cause inflammation of the tongue, known as glossitis. The tongue may appear swollen, smooth, and red.

- **Loss of Appetite and Weight Loss:** Some individuals with megaloblastic anemia may experience a decrease in appetite, leading to unintended weight loss.
- **Neurological Symptoms:** In severe cases of vitamin B12 deficiency, neurological symptoms can occur. These may include numbness or tingling in the hands and feet (peripheral neuropathy), coordination problems, memory loss, confusion, and even depression.

DIAGNOSIS:

- **Family history & Medical history**
- **Physical examination:** The healthcare provider will check for signs & symptoms.
- **Complete Blood Count (CBC):** A CBC provides an overview of the blood's cellular components. In megaloblastic anemia, the CBC may reveal a low red blood cell count (anemia), an increased mean corpuscular volume (MCV) indicating the presence of larger red blood cells (macrocytosis), and a decreased hemoglobin and hematocrit level.
- **Peripheral Blood Smear:** A blood smear involves examining a small sample of blood under a microscope. In megaloblastic anemia, the smear may show the presence of larger and immature red blood cells (megaloblasts) with characteristic features, such as a high nuclear-to-cytoplasm ratio.
- **Vitamin B12 and Folate Levels:** Blood tests can measure the levels of vitamin B12 and folate in the bloodstream. Low levels of vitamin B12 or folate can indicate a deficiency contributing to megaloblastic anemia.
- **Intrinsic Factor Antibody Test:** This test is specifically performed to check for pernicious anemia, an autoimmune condition that affects vitamin

B12 absorption. It detects the presence of antibodies against intrinsic factor, a protein required for vitamin B12 absorption.

- **Schilling Test:** The Schilling test is conducted to assess the absorption of vitamin B12. It involves giving a small dose of radioactive vitamin B12 and measuring the amount excreted in the urine. This test can help determine if there is a problem with vitamin B12 absorption.
- **Bone Marrow Examination:** In some cases, a bone marrow biopsy may be performed to examine the cells in the bone marrow directly. This can provide more detailed information about the presence of megaloblasts and help confirm the diagnosis.

PHARMACOLOGICAL MANAGERMENTS:

Vitamin B12 (Cobalamin) Supplements:

For megaloblastic anemia caused by vitamin B12 deficiency, the primary treatment is vitamin B12 supplementation. It can be administered in different forms, including oral tablets, sublingual tablets, intramuscular injections, or nasal sprays. Examples of vitamin B12 supplements include:

- **Cyanocobalamin**
- **Hydroxocobalamin**
- **Methylcobalamin**

Folate (Vitamin B9) Supplements:

In cases where megaloblastic anemia is caused by folate deficiency, folate supplementation is the main treatment. Folate supplements are usually available in oral tablet form. Examples of folate supplements include:

- **Folic Acid**

NON PHARMACOLOGICAL MANAGERMENTS:

- **Balanced Diet:** Maintain a well-balanced diet that includes foods rich in vitamin B12 and folate. Good dietary sources of vitamin B12 include meat,

poultry, fish, eggs, dairy products, and fortified cereals. Foods rich in folate include leafy green vegetables, legumes, citrus fruits, and fortified grains.

- **Vitamin Supplements:** If you have a known deficiency or are at risk of megaloblastic anemia, consider taking vitamin supplements under the guidance of a healthcare professional. This is especially important for individuals following restrictive diets, such as strict vegetarians or vegans who may need vitamin B12 supplementation.
- **Prenatal Care:** Pregnant women should receive adequate prenatal care, including regular check-ups and nutritional counseling. Folate requirements are significantly increased during pregnancy, and supplementation is often recommended to prevent neural tube defects in infants.
- **Treat Underlying Conditions:** Properly manage underlying conditions that may lead to megaloblastic anemia, such as pernicious anemia, gastrointestinal disorders, or malabsorption syndromes. This may involve medication, dietary adjustments, or other medical interventions as advised by healthcare professionals.
- **Medication Management:** If you are taking medications that can interfere with vitamin B12 or folate absorption or metabolism, discuss with your healthcare provider about potential supplementation or monitoring to prevent deficiencies.
- **Regular Check-ups:** Regularly visit your healthcare provider for routine check-ups and screenings to detect any potential vitamin deficiencies or underlying conditions early on.