

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/339899087>

Yield maximization through higher fertilization and lodging management in irrigated timely sown wheat (*Triticum aestivum* L.) under eastern sub-Himalayan plains

Article · December 2020

DOI: 10.22271/09746315.2020.v16.i3.1384

CITATIONS

2

READS

229

4 authors:



Triptesh Mondal

Centurion University of Technology and Management

27 PUBLICATIONS 57 CITATIONS

[SEE PROFILE](#)



Biplab Mitra

Uttar Banga Krishi Vishwavidyalaya

44 PUBLICATIONS 276 CITATIONS

[SEE PROFILE](#)



Arnab Roy Chowdhury

Bihar Agriculture University

18 PUBLICATIONS 29 CITATIONS

[SEE PROFILE](#)



Saikat Das

Uttar Banga Krishi Vishwavidyalaya

19 PUBLICATIONS 55 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



AICRP on Weed Management [View project](#)



AICRP on Wheat and Barley, UBKV Centre formulated from ICAR-IIWBR, Karnal [View project](#)



Yield maximization through higher fertilization and lodging management in irrigated timely sown wheat (*Triticum aestivum* L.) under eastern sub-Himalayan plains

T. MONDAL, *B. MITRA, A. ROY CHOWDHURY AND ¹S. DAS

Department of Agronomy, ¹All India Coordinated Wheat and Barley Improvement Project
Uttar Banga Krishi Viswavidyalaya, Pundibari, Cooch Behar - 736165, West Bengal

Received : 30.09.2020 ; Revised : 13.11.2020 ; Accepted : 15.11.2020

DOI : <https://doi.org/10.22271/09746315.2020.v16.i3.1384>

ABSTRACT

An experiment was conducted during rabi season of 2016-17 and 2017-18 under eastern sub-Himalayan plains of West Bengal, India with the objective to maximize grain yield of wheat (*Triticum aestivum* L.) through higher fertilization and lodging management. The design of the experiment was split-plot with 4 nutrient management options viz. No application of fertilizer (F₀); Recommended dose (RD) (150, 60, 40 kg N, P₂O₅, K₂O ha⁻¹) (F₁); 150% RD (225, 90, 60 kg N, P₂O₅, K₂O kg ha⁻¹) (F₂); 150% RD (225, 90, 60 N, P₂O₅, K₂O kg ha⁻¹) + 15 t ha⁻¹ FYM (F₃) in main-plots and 4 growth regulator spraying options viz. No spraying of chemicals, spraying of water twice (G₀); spraying of chlormequat chloride twice (G₁); spraying of tebuconazole twice (G₂); spraying of chlormequat chloride + tebuconazole twice (G₃) in sub-plots each replicated thrice. It was revealed that 150% RD + 15 t ha⁻¹ FYM treatment recorded the highest grain yield (5.61 t ha⁻¹) followed by 150% RD only (5.30 t ha⁻¹). Chlormequat chloride (0.2%) + tebuconazole (0.1%) spray at 1st node and flag leaf development stage provided lowest lodging score (1.65) reflecting its reliability in lodging management. Thus 150% RD with lodging management through chlormequat chloride (0.2%) + tebuconazole (0.1%) spraying twice at 1st node and flag leaf stage indicated its superiority in terms of benefit:cost ratio (B:C) (2.51:1) with higher agronomic N use efficiency (ANUE) (23.84 kg grain kg N⁻¹) and economic N use efficiency (ENUE) (2.80 kg grain invested in N⁻¹).

Keywords: Chlormequat chloride, fertilization, lodging management, wheat and tebuconazole

The yield of wheat (*Triticum aestivum* L.), the most important rabi food grain crop of India, has started stagnating or even declining from mid-eighties. There has been a wide gap between the target and actual wheat production of the country (Pathak *et al.*, 2003). The potential yield of irrigated wheat is about 5-7 t ha⁻¹ in the state of West Bengal but the actual yield obtained is much lower, thereby indicating a scope of its improvement through appropriate crop management practices (Bhattacharyya *et al.*, 2008). Nitrogen (N) is the key nutrient behind this yield reduction which actually controls the crop development in cereal based cropping systems (Samborski *et al.*, 2009). Phosphorus (P) is also an important plant nutrient next only to N and classed as a major plant nutrient along with N and potassium (K). Farm yard manure (FYM) incorporation along with inorganic nutrients considerably increased all the growth and yield attributing parameters along with yield and nutrient uptake *vis-à-vis* highest nutrient recovery (Bhaduri and Gautam, 2012).

Lodging, the wind induced permanent displacement of cereal stems from the vertical, can cut the profitability of a cereal crop drastically through lower grain yield (Sterling *et al.*, 2003). Lodging was induced with increasing N fertility beyond certain limit which in turn decreased grain yield and its components (Fischer and

Stapper, 1987). After stem lodging, the cells on the lower side of the node elongate and force the stem erect. As plants mature, the stem cells force and are no longer capable of elongation to enable plant recovery (Telkar *et al.*, 2017). Plant growth regulators (PGRs) are often used to improve lodging traits, protect yield potentials and restrain elongation of internodes which are associated with reduction of plant height (Zhang *et al.*, 2017). It is well perceived that for harnessing full potential of genetic makeup higher dose of fertilization is adopted, however, at the same time appropriate lodging management is essential as plants are susceptible to lodging with increased fertilization. With this backdrop, a two-year investigation was undertaken to assess the effects of higher level of nutrition towards yield maximization and to manage lodging due to higher doses of fertilizer application in timely sown, irrigated wheat.

MATERIALS AND METHODS

Field trials were conducted during two winter seasons in 2016-17 and 2017-18 at the Instructional Farm of Uttar Banga Krishi Viswavidyalaya (UBKV), Pundibari, Cooch Behar, West Bengal under eastern sub-Himalayan plains located at 26°24'02.1"N latitude, 89°23'21.5"E longitude and at 43 m above mean sea level (MSL). The experimental soil (0-15 cm soil layer) was sandy loam in texture with pH 5.51, soil organic carbon (C) 0.79%,

*Email: bipmitra@yahoo.com

mineralizable N 175.70 kg ha⁻¹, Bray I P 23.30 kg ha⁻¹ and NH₄OAc extractable K 86.60 kg ha⁻¹.

The experiment was laid out in split plot design with three replications. Four levels of nutrition viz. No application of fertilizer (F₀); Recommended dose (RD) (150, 60, 40 kg N, P₂O₅, K₂O ha⁻¹) (F₁); 150% RD (225, 90, 60 kg N, P₂O₅, K₂O kg ha⁻¹) (F₂); 150% RD (225, 90, 60 N, P₂O₅, K₂O kg ha⁻¹) + 15 t ha⁻¹ FYM (F₃) were randomly allotted in the main-plots and four treatments consisting of various lodging arresting chemicals viz. no spraying of chemicals, spraying of water twice (G₀); spraying of chlormequat chloride (0.2%) twice at first node and flag leaf stage (G₁); spraying of tebuconazole (0.1%) twice at first node and flag leaf stage (G₂); spraying of chlormequat chloride (0.2%) + tebuconazole (0.1%) twice at first node and flag leaf stage (G₃) were randomly allocated in the sub-plots. The one-third of nitrogen fertilizers was applied as basal along

with full P and half K fertilizers. The remaining N was applied in two equal splits, once 20-22 days after sowing (DAS) and other at 40-42 DAS with remaining half K. The individual sub-plot size was 8 × 1.8 m (14.4 m²). Wheat was cultivated after puddled transplanted rice in both the years of experiment. Wheat variety HD 2967 was used in this experiment. Seeds were sown in rows 20 cm apart with a seed rate of 100 kg ha⁻¹ during November 18 and 20 in first year and second year, respectively. Nutrients were applied as per treatment. In the treatment consisting of FYM, it was applied 15 days before the final land preparation. Among the micronutrients, boron (B) was applied @0.2% with Solubor (B 20%), once at 35-40 days after sowing (DAS) and the subsequent one at 55-60 DAS and zinc (Zn) was applied with B in the 2nd spray, i.e., @0.1% with chelated Zn (Zn-EDTA 12%). Lodging score at harvest was calculated by the following formula (Fischer and Stapper, 1987).

$$\text{Lodging score} = \frac{(\% \text{ plot area lodged} \times \text{angle of lodging from the vertical})}{90}$$

Agronomic nitrogen use efficiency (ANUE) and economic nitrogen use efficiency (ENUE) of wheat were

calculated by the following formula of Nyborg et al. (1995) and Ahmad *et al.* (2012), respectively.

$$\text{ANUE} = \frac{(\text{Grain yield with given N} - \text{Grain yield without N})}{\text{N application rate}} \text{ kg grain kg N}^{-1}$$

$$\text{ENUE} = \frac{(\text{Grain yield with given N} - \text{Grain yield without N})}{\text{Value of N application rate}} \text{ kg grain ? invested in N}^{-1}$$

Wheat was harvested manually from the ground level by sickle. Net plot area was harvested skipping the border rows approximately 7-10 days after physiological maturity for estimating grain and straw yield of wheat. Gross return, net return and benefit-cost (B:C) ratio were calculated by taking the prevailing market price of wheat grain (Rs. 16.00 kg⁻¹) and wheat straw (Rs. 1.50 kg⁻¹) during that period. Data followed the homogeneity test and pooling was done over two years. Statistical analysis of data was done with the help of SPSS software.

RESULTS AND DISCUSSION

Growth attributes

Plant height at harvest was increased by 5.43% in 150% of recommended NPK treatment over the treatment receiving only recommended dose (RD). Lodging arresting chemicals had significant effect on plant height and two sprays of chlormequat chloride (0.2%) + tebuconazole (0.1%) at first node and flag leaf stage showed lowest plant height (81.44 cm) among all the PGR spraying options (Table 1). In case of plant height at harvest, significant interaction effect between fertilization rates and growth regulator spraying options

was recorded. It was also revealed that fertilization significantly enhanced tiller and biomass production. Highest tiller density was achieved in combined application of 150% RD and 15 t ha⁻¹ FYM (466 m⁻² at 90 DAS) and lowest tiller density was observed in no fertilizer treated plots (208 m⁻² at 90 DAS) (Table 1). Higher N rate was mainly responsible for the enhancement in tiller production. This finding is in agreement with the finding of Balkcom and Burmester (2015). Grain + straw biomass (t ha⁻¹) was enhanced by 18.56-32.40% in NPK-rich plots over 100% RD plots. It was mainly due to N, being an important constituent of amino acids, proteins and protoplasm, which highly influenced grain + straw biomass through increase in shoot number, cell expansion and better photosynthate utilization under more N availability (Kumar *et al.*, 2017). Leaf area index (LAI) at 90 days after sowing (DAS) was significantly increased due to higher NPK rates over recommended rate of NPK and it was increased by 23.11-30.41%. However, the PGR spray had no significant effect on LAI of wheat at anthesis. Toyota *et al.* (2010) reported similar observations about chlormequat chloride (0.2%) spray in wheat.

Physiological maturity and lodging score at harvest

Physiological maturity dates were studied with respect to both fertilization and foliar application of growth regulators. There was significant increase in the duration for attaining the physiological maturity with higher doses of nutrient application than no NPK applied treatment (Table 1). Similar observations were reported by Mondal *et al.* (2018). This was mainly because of sufficient distribution of N to flag leaf in N-rich treatments which increased stay-green duration (Gaju *et al.*, 2014) while plants attended forced maturity under N starvation. It was observed that higher lodging in wheat occurred with 225 kg N ha⁻¹, previously reported by Hobbs *et al.* (1998). Higher NPK treated plots showed

significantly higher lodging scores (4.29 in 150% recommended NPK and 7.54 in 150% recommended NPK with 15 t ha⁻¹ FYM treatment) than the fertilized plots with recommended dose of NPK (1.40 in recommended NPK treatment). No lodging was recorded in the plots having no fertilizer application. Growth regulators had significant effect on lodging score. Least lodging score (1.60) was estimated from the treatment where combined spraying of 0.2% chlormequat chloride and 0.1% tebuconazole was adopted twice at 1st node and flag leaf stage of wheat. In case of lodging score at harvest, interaction effect was found significant between fertilization treatments and lodging arresting chemical treatments (Table 1).

Table 1: Growth attributes, physiological maturity and lodging score of wheat as affected by fertilization and growth regulator (mean over 2 years)

Treatments	Growth attributes				Physiological maturity (DAS)	Lodging score* at harvest
	Plant height at harvest (cm)	Tiller density m ⁻² at 90 DAS	LAI at 90 DAS	Grain + Straw biomass (t ha ⁻¹)		
Fertilization (F)						
F ₀	67.52	208	1.64	4.38	120.00	0.00
F ₁	90.13	384	4.11	9.32	124.50	1.40
F ₂	95.02	429	5.06	11.05	125.88	4.29
F ₃	92.97	466	5.36	12.34	126.96	7.54
SEm (±)	0.65	6.28	0.12	0.20	0.41	0.57
LSD (0.05)	4.41	32.78	0.81	1.42	1.74	3.28
Growth regulator (GR) spraying						
G ₀	90.72	363	3.92	9.20	124.04	5.84
G ₁	84.23	369	3.97	9.06	124.29	2.29
G ₂	89.25	373	4.02	9.19	124.42	3.44
G ₃	81.44	383	4.27	9.64	124.58	1.65
SEm (±)	0.63	4.24	0.17	0.12	0.31	0.40
LSD (0.05)	2.68	NS	NS	NS	NS	2.37
F × GR.SEm (±)	1.30	12.57	0.25	0.40	0.82	1.03
LSD (0.05)	5.36	NS	NS	NS	NS	4.74
GR × FS.SEm (±)	1.13	13.25	0.38	0.65	0.92	1.26
LSD (0.05)	4.62	NS	NS	NS	NS	3.96

Note: DAS: Days after sowing, *Angle of stem lodging was 45°

Yield attributes and yield

Nutrient management options had significant effect on yield attributes *viz.*, number of spike m⁻², number of filled grain spike⁻¹, spike length and 1000-grain weight as previously reported by Mondal *et al.* (2018). Highest number of spike m⁻² (339 m⁻²) was noted in treatments with 225, 90, 60 kg N, P₂O₅, K₂O kg ha⁻¹ + 15 t ha⁻¹ FYM (Table 2). Application of higher N rate increased effective tiller production rate (Yang *et al.*, 2019) which ultimately increased the spike density in wheat. Although

150% recommended NPK application recorded the highest spike length (10.59 cm) but highest number of filled grain spike⁻¹ was recorded from 150% recommended NPK + 15 t ha⁻¹ FYM treatment (43.58). This increment in number of filled grain spike⁻¹ was due to combined application of organic manure and inorganic fertilizers as reported previously by Singh *et al.* (2011). Foliar application of growth regulators also imposed significant influence on spike length. It was revealed that 1000-grain weight (42.48 g) was recorded highest

from 150% RD treatment. This was due to higher N application rate as reported previously by Krylov and Pavlov (1989). Fertilization significantly influenced grain yield, straw yield and harvest index. Highest grain yield was obtained under 150% RD along with 15 t ha⁻¹ FYM treatments (5.61 t ha⁻¹). This was also due to combined application of organic manure and inorganic fertilizers which resulted in higher number of yield attributes under a better nutritional environment. The findings were in conformity with the report of Singh *et al.* (2011). Straw yield followed similar trend of grain yield and was recorded highest under 150% RD + 15 t ha⁻¹ FYM treatment (7.34 t ha⁻¹) with maximum harvest index (45.49%) under 150% RD treatment.

ANUE, ENUE and production economics

The ANUE and ENUE values were lower in N-rich plots, similar with the previous report of Mondal *et al.* (2018). Highest ANUE values were recorded in recommended N based treatments as compared to excess N applied treatments. The trend was similar for ENUE

also (Table 3). The data revealed that there was yield enhancement under higher N doses; however, it was not a proportionate enhancement as reflected by ANUE and ENUE values. Excess N application under 150% RD treatments, particularly during initial stages, may not always put the plant to use it efficiently. That is why the ANUE *vis-à-vis* ENUE did not result in enhanced values over 100% RD treatment, rather it reflected its misuse through which there was higher loss of N. Under recommended N treatments, the N rate of application was much lower than the treatments comprising 150% RD and in this case N was effectively utilized by the wheat crop. That is why plant nitrogen use efficiency including ANUE and ENUE was gradually decreased due to over application of N (Yang *et al.*, 2019). Mondal *et al.* (2018) and Mitra *et al.* (2019) also reported higher N use efficiencies with lower or recommended N levels in wheat at eastern sub-Himalayan plains as compared to N rich plots which actually reflected the need of site specific nutrient management.

Table 2: Yield attributes, grain yield, straw yield and harvest index of wheat as affected by fertilization and growth regulator (mean over 2 years)

Treatments	Yield attributes				Grain yield (t ha ⁻¹)	Straw yield (t ha ⁻¹)	Harvest index (%)
	No. of spike m ⁻²	Spike length (cm)	No. of filled grain spike ⁻¹	Test weight (g)			
Fertilization (F)							
F ₀	128	7.01	34.17	36.50	1.81	3.14	37.00
F ₁	269	10.27	42.92	41.95	4.27	5.64	43.08
F ₂	318	10.59	43.13	42.48	5.30	6.35	45.49
F ₃	339	10.23	43.58	42.05	5.61	7.34	43.32
SEm (±)	5.10	0.048	1.09	0.27	0.13	0.17	1.157
LSD (0.05)	32.78	0.39	5.30	1.04	0.91	0.96	4.58
Growth regulator (GR) spraying							
G ₀	259	9.70	40.50	40.52	4.14	5.62	42.42
G ₁	260	9.39	40.38	40.60	4.14	5.52	42.86
G ₂	260	9.67	41.25	40.91	4.25	5.57	43.28
G ₃	274	9.34	41.67	40.95	4.46	5.77	43.60
SEm (±)	4.07	0.04	1.54	0.37	0.13	0.25	1.43
LSD (0.05)	NS	0.21	NS	NS	NS	NS	NS
F × GR S.E_m (±)	10.28	0.07	2.38	0.54	0.26	0.34	2.59
LSD (0.05)	NS	0.39	NS	NS	NS	NS	NS
GR × F S.E_m (±)	12.34	0.08	2.59	0.72	0.24	0.50	2.90
LSD (0.05)	NS	0.45	NS	NS	NS	NS	NS

The data on production economics revealed the superiority of the treatments consisting of 150% RDF. The highest net return (Rs. 60693 ha⁻¹) and benefit-cost ratio (2.51:1) were recorded from 150% RDF with chlormequat chloride (0.2%) + tebuconazole (0.1%)

spraying treatment combination (Table 3). Highest net return and benefit-cost ratio reflected the reliability of this treatment combination towards achieving higher profitability. Although grain yield was highest under 150% RDF + 15 t ha⁻¹ FYM treated plots but the benefit-

cost ratio were very poor in 150% RDF + 15 t ha⁻¹FYM-chlormequat chloride (0.2%) + tebuconazole (0.1%) spraying treatment combination reflecting that the extra

cost incurred for FYM was much higher than its proportionate return. However, FYM had a carry-over effect on subsequent crop in the rotation with better soil health.

Table 3: ANUE, ENUE and economics (net returns and benefit cost ratio) of wheat as affected by fertilization - growth regulator spraying treatment combinations (mean of 2 years)

Treatment combinations	ANUE* (kg grain kg N ⁻¹)	ENUE** (kg grain Rs. invested in N ⁻¹)	Net return (Rs. ha ⁻¹)	B:C
F ₀ -G ₀	-	-	5563	1.20:1
F ₀ -G ₁	-	-	3523	1.12:1
F ₀ -G ₂	-	-	3695	1.13:1
F ₀ -G ₃	-	-	3767	1.12:1
F ₁ -G ₀	27.29	3.03	39179	2.13:1
F ₁ -G ₁	27.58	3.06	38107	2.05:1
F ₁ -G ₂	28.88	3.21	42076	2.19:1
F ₁ -G ₃	30.21	3.36	43957	2.18:1
F ₂ -G ₀	20.79	2.56	54144	2.42:1
F ₂ -G ₁	20.98	2.45	48921	2.23:1
F ₂ -G ₂	22.96	2.65	56422	2.46:1
F ₂ -G ₃	23.84	2.80	60693	2.51:1
F ₃ -G ₀	18.50	0.27	30630	1.45:1
F ₃ -G ₁	18.56	0.27	31774	1.46:1
F ₃ -G ₂	18.84	0.27	30417	1.44:1
F ₃ -G ₃	19.60	0.28	33063	1.47:1

Note: *FYM contains 0.40% N, 0.20% P₂O₅ and 0.18% K₂O (after chemical analysis of FYM); **Rs. invested per kg of N were 19.57 for RDF, 150% RDF and 72.21 for 150% RDF + 15 t ha⁻¹ FYM

The study unveiled that increased fertilization through 150% of recommended NPK not only increased grain yield of wheat but also resulted in higher economic benefit. Inclusion of FYM @ 15 t ha⁻¹ along with 150% of recommended NPK may increase the yields further, but may not be economic. However, higher N application recorded lower values of ANUE and ENUE indicating the wastage of nitrogen. Among the lodging arresting chemical treatments, spraying of chlormequat chloride @0.2% + tebuconazole @0.1% showed good control of lodging. Therefore, 150% recommended rate of NPK through chemical fertilizers along with combined spraying of chlormequat chloride (0.2%) and tebuconazole (0.1%) twice at 1st node and flag leaf stage of wheat could maximize grain yield in wheat with better control of lodging and increased profitability in eastern sub-Himalayan plains.

ACKNOWLEDGEMENT

Contributions of All India Coordinated Wheat and Barley Improvement Project (AICW & BIP), Coochbehar Centre, UBKV for providing necessary financial and technical support towards successful conduction of the field trial are duly acknowledged.

REFERENCES

- Ahmed, S., Ali, H., Ismail, M., Shahzad, M.I., Nadeem, M., Anjum, M.A., Zia-Ul-Haq, M., Firdous, N. and Khan, M.A. 2012. Radiation and nitrogen use efficiencies of C3 winter cereals to nitrogen split application. *Pakistan J. Bot.*, **44**(1): 139-149.
- Balkcom, K.S. and Burmester, C.H. 2015. Nitrogen applications for wheat production across tillage systems in Alabama. *Agron. J.*, **107**(2): 425-434.
- Bhaduri, D. and Gautam, P. 2012. Balanced use of fertilizers and FYM to enhance nutrient recovery and productivity of wheat (*Triticumaestivum* cv. UP-2382) in a Mollisol of Uttarakhand. *Int. J. Agric., Environ. Biotechnol.*, **5**(4): 435-439.
- Bhattacharyya, P., Mallick, E.H., Pathak, H. and Ladha, J.K. 2008. Wheat in West Bengal. Directorate of Agriculture, West Bengal and Rice-Wheat Consortium for Indo-Gangetic Plains, New Delhi, India, pp. 35.
- Donald, C.M. and Hamblin, J. 1976. The biological yield and harvest index of cereals as agronomic and plant breeding criteria. *Adv. Agron.*, **29**: 361-405.
- Fischer, R.A. and Stapper, M. 1987. Lodging effects on high yielding crops of irrigated semi-dwarf wheat. *Field Crops Res.*, **17**: 245-258.

- Gaju, O., Allard, V., Martre, P., Gouis, J.L., Moreau, D., Bogard, M.S., Hubbart, M. and Foulkes, M.J. 2014. Nitrogen partitioning and remobilization in relation to leaf senescence, grain yield and grain nitrogen concentration in wheat cultivars. *Field Crops Res.*, **155**: 213-223.
- Hobbs, P.R., Sayre, K.D. and Ortiz-Monasterio, I. 1998. Increasing wheat yields through agronomic means. In: Nagarajan, S., Singh, G. and Tyagi, B.S. (Eds.), *Wheat Research Beyond 2000 AD*. Proceedings of the International Group Meeting on Research Needs Beyond 2000 AD, Karnal (India). 12-24 Aug, 1997, Narosa Publishing House, New Delhi, India.
- Krylov, Y.A.I. and Pavlov, V.D. 1989. Effect of fertilizer on yield and protein contents in wheat grain. *Agrochimiya*, **1**: 49-51.
- Kumar, V., Raghuvanshi, N. and Singh, R.A. 2017. Effect of different combination of FYM and urea on growth and yield of wheat (*Triticum aestivum* L.). *Bull. Env.Pharma. Life Sci.* (special issue), **6**(2): 395-398.
- Mitra, B., Majumdar, K., Dutta, S. K., Mondal, T., Das, S., Banerjee, H., Ray, K. and Satyanarayana, T. 2019. Nutrient management in wheat production system under conventional and zero tillage in eastern sub-Himalayan plains of India. *Ind. J. Agril. Sci.*, **89**(5): 775-784.
- Mondal, T., Mitra, B. and Das, S. 2018. Precision nutrient management in wheat (*Triticum aestivum*) using Nutrient Expert®: Growth phenology, yield, nitrogen use efficiency and profitability under eastern sub-Himalayan plains. *Indian J. Agron.*, **63**(2): 174-180.
- Nyborg, M., Solberg, E.D., Izaurralde, R.C., Malhi, S.S. and Molina-Ayala, M. 1995. Influence of long term tillage, straw and fertilizer on barley yield, plant N uptake and soil-N balance. *Soil Till. Res.*, **36**: 165-174.
- Pathak, H., Aggarwal, P.K., Roetter, R., Kalra, N., Bandyopadhyaya, S.K., Prasad, S. and Keulen, H.V. 2003. Modeling the quantitative evaluation of soil nutrient supply, nutrient use efficiency and fertilizer requirements of wheat in India. *Nut.Cycl.Ag.*, **65**: 105-113.
- Samborski, S.M., Tremblay, N. and Fallon, E. 2009. Strategies to make use of plant sensors-based diagnostic information for nitrogen recommendations. *Agron. J.*, **101**(4): 800-816.
- Singh, C.M., Sharma, P.K., Kishor, P., Mishra, P.K., Singh, A.P., Verma, R. and Raha, P. 2011. Impact of integrated nutrient management on growth, yield and nutrient uptake by wheat (*Triticum aestivum* L.). *Asian J. Agric. Res.*, **5**(1): 76-82.
- Sterling, S., Baker, C.J., Berry, P.M. and Wade, A. 2003. An experimental investigation of the lodging of wheat. *Agric. For. Met.*, **119**: 149-165.
- Telkar, S.G., Solanki, S.P.S., Chouhan, S., Kumar, R. and Nikas, S.B. 2017. Crop Lodging on Cereals: Causes, Effect and Control. *Biomolecule Reports - An Int. e-Newslet.*, pp: 1-5.
- Toyota, M., Shiotsu, F., Bian, J., Morokuma, M. and Kusutani, A. 2010. Effects of reduction in plant height induced by chlormequat on radiation interception and radiation-use efficiency in wheat in southwest Japan. *Plant Prod. Sci.*, **13**(1): 67-73.
- Yang, D., Cai, T., Luo, Y. and Wang, Z. 2019. Optimizing plant density and nitrogen application to manipulate tiller growth and increase grain yield and nitrogen-use efficiency in winter wheat. *Peer J.*, pp. 1-26.
- Zhang, Y., Su, S., Tabori, M., Yu, J., Chabot, D., Baninasab, B., Wang, X., Ma, B.L., Li, C. and Khanizadeh, S. 2017. Effect of selected plant growth regulators on yield and stem height of spring wheat in Ontario. *J. Agric. Sci.*, **9**(12): 30-42.