

PLANT GENOMICS

Thomas Blake, Victoria Carollo Blake and Jackie Campbell

Department of Plant Sciences and Plant Pathology, Montana State University, USA

Keywords: DNA, gene expression, gene regulation, genome, mapping, RNA, siRNA, transposon.

Contents

1. Introduction to Genomics
 2. History of Plant Genomics
 3. Structure and Composition of Plant Genomes
 4. Mapping Plant Genomes
 - 4.1. Mapping Technologies
 - 4.2. The Way Forward
 - 4.3. Mapping Important Genes
 - 4.4. Classic Quantitative Trait Locus (QTL) Analysis
 - 4.5. Association Analysis and Germplasm Utilization
 - 4.6. Positional Cloning: From Map Location to Gene
 5. Functional Genomics
 - 5.1. Northern Blot
 - 5.2. Electronic Northern Blots
 - 5.3. SAGE
 - 5.4. Real Time PCR (rtPCR)
 - 5.5. DNA Microarrays
 - 5.6. Affymetrix Microarrays
 - 5.7. Data Repositories
 6. Small RNA Molecules
 - 6.1. RNA Interference (RNAi)
 - 6.2. Cross Protection
 - 6.3. Virus Induced Gene Silencing
 7. Conclusion: The Future of Plant Genomics
- Glossary
Bibliography
Biographical Sketches

Summary

Plant genomes dramatically vary in size and in function. In years past, genome size rather than economic or environmental utility determined which plant species were best characterized at the level of DNA sequence. Still, the broad outlines of plant genome organization and evolution were well-understood for most important crop species by the mid-1970's.

The diversity, both within and among species, that modern genetic tools reveal, is proving to be dazzling, and plant diversity is now being explored and utilized more effectively than ever before.

Plant biologists uncovered many of the most interesting facets of eukaryotic genomics, including transposition, retro-transposition and small RNA mediated gene regulation. New tools that are permitting the exploration, mapping and remodeling of plant genomes hold the promise to enable the consistent gains in plant productivity that our growing population and declining environmental resources will demand in the 21st century.

1. Introduction to Genomics

The focus of this chapter is on the plant nuclear genome. Plant chloroplasts and mitochondria maintain genomes, and plant mitochondrial genomes are large, variable and enormously interesting. Nuclear genomes of plants have been sculpted by inter-specific hybridization, polyploidy, transposition, retro-transposition, translocation and illegitimate recombination events. Wheat and maize provide excellent examples of several of these processes.

Durum wheat, used to make pasta, resulted from the hybridization of two diploid wheat grass species, *Triticum monococcum* and a grass very much like *Triticum speltoides*. The nuclear genomes of each of these related grasses contain fourteen chromosomes, and occasional failures of meiosis result in pollen and egg cells with fourteen, rather than the expected seven chromosomes. Durum wheat resulted from the fusion of two of these unreduced gametes, resulting in a plant with 28 chromosomes. Bread wheat resulted from the hybridization of an unreduced durum wheat egg by an unreduced pollen grain from the related grass, *Triticum tauschii*. Durum and bread wheat are examples of polyploids, organisms with multiple complete chromosome sets, derived from inter-specific hybridization events.

Maize achieves much of its remarkable diversity through the activity of transposons. These are DNA sequences that contain specific repeated sequences at their left and right boundaries that are recognized by enzymes that catalyze their movement from place to place in maize chromosomes. Transposons and retro-transposons help move genes and parts of genes throughout the genomes of higher plants, generating diversity, and also generate mutations by moving within or near genes. During meiosis, chromosomes occasionally mispair. When recombination occurs between mispaired chromosomes, part of one chromosome is swapped with part of another. One translocation differentiates the genomes of wheat and barley.

Gene duplication and divergence is common, and all of these genetic mechanisms result in what has been termed emergent complexity. We heterotrophs (organisms that derive nutrition by consuming other organisms) benefit from the variation and complexity that plant genome diversity provides. Human motivation for studying plant genomes is simple. Plants make our lives possible, and their beauty and utility are determined by the genetic programs stored within their genomes.

We all depend on plants for the air we breathe and the food we eat. As we move through the 21st century, our dependence on the plants that service our environment and provide food, fuel and fiber will become more evident. Our ability to utilize the genetic resources provided by evolution and to engineer novel solutions to current and coming

problems will be tested. Plant genomics is becoming an invaluable tool to those involved in crop improvement and germplasm management.

Plant nuclear genomes range in size from less than 100 million base pairs to more than 100 billion base pairs. Several explanations were offered to explain the apparent lack of correlation between genome size and plant form and function. The best current explanation for this variation is that replicative mobile genetic elements, primarily retro-transposons, invade eukaryotic chromosomes, and that genome size variation is primarily a consequence of different species' resistance to or tolerance of thousands of copies of these virus-like DNA sequences.

Genetics, bioinformatics, physiology and descriptive biology all contribute to genomics. Several plant genomes have been completely sequenced, but DNA sequence alone does not provide complete insight into the genomes' dynamic nature. We are now discovering that even among humans, substantial variation in the number of copies of genes can be observed among different individuals. Many of our most important crop plants (e.g. wheat and soybean) originated through inter-specific hybridization or whole genome duplication events, suggesting that we can expect plant genome architecture to be characterized by even greater variation both within and among species.

2. History of Plant Genomics

Formal research into the function of the plant genome started with Mendel in 1866. In a remarkable paper, Mendel demonstrated the particulate nature of genes, and provided proof of both segregation and independent assortment of genes. A surge of interest in genetics and its applications followed the confirmation, popularization and extension of Mendel's work (see also: *Plant Breeding and Genetics*).

In 1987 McClintock demonstrated the surprising capacity for specific maize genotypes to generate both forward and reverse mutations within several well-characterized genes at remarkably high rates. Her groundbreaking work with maize transposable elements led to an understanding of the phenomenology of transposition in both pro- and eukaryotes. Her insight into the adaptive role of transposition in the development of variation that contributes to local adaptation remains a central contribution to modern biology.

Several technologies ranging from quantitative Fuelgen stain analysis to DNA re-naturation analysis (see below) indicated that most crop genomes were large, and were full of repeated sequences. The activity of transposons and retro-transposons helped explain the initially confusing picture of higher plant genomes provided by DNA re-naturation analysis. These mobile genetic elements share several features with viruses, and have been termed genetic parasites. Still, they help generate the variation that natural selection acts upon, and in doing so contribute to plants' amazing capacity to adapt.

One of the mysteries of plant biology, the obvious improved performance of hybrid maize plants when contrasted with their parents, was initially correctly interpreted by Jones (1917), as a consequence of genetic linkage between undesirable recessive alleles

and desirable dominant alleles. Maize breeders exploit this feature of the maize genome by making crosses between appropriate parents, and producing hybrids that are homozygous for as few undesirable recessive alleles as possible. When Sax demonstrated in the 1920s that a gene impacting seed size in *Phaseolus vulgaris* was linked to a gene conferring seed pigmentation, scientists worldwide recognized the potential value of genome-wide linkage maps, which relate the relative positions of genes or markers on a chromosome based on the frequency of crossover between them during meiosis.

With enough genetic markers, one could utilize linkage analysis to identify all the chromosomal locations of the genes impacting plant performance, and could, theoretically, develop the 'best' variety for any environment. Elegant cytogenetic tools were developed to permit efficient chromosomal location of genes controlling Mendelian characters. By 1970, maize, wheat, pea and barley all had well-populated linkage maps. The typically recessive morphology-impacting genes that had been mapped were identified in specific genotypes, and compiling a sufficient number of these rare genetic variant alleles into a genetic tester line typically resulted in plants that performed poorly.

Botstein *et al.* (1980) pointed to DNA sequence polymorphism analysis as an approach that could, potentially, provide enough genetic markers for humans to permit genome-wide linkage map construction. Restriction fragment length polymorphisms (RFLPs) provided endogenous genetic markers sufficiently informative to make genome-wide linkage map construction feasible. For examples of the RFLP-based Southern Blots that permitted linkage map construction in barley, wheat and oat, go to <http://wheat.pw.usda.gov>. Linkage maps produced by RFLP analysis are also viewable on GrainGenes. Plant geneticists and plant breeders adopted RFLP analysis, resulting in the development of excellent, reasonably dense linkage maps in a wide array of plant species without the performance decay observed in plants carrying mutations that impacted plant phenotype (Tanksley and Fulton, 2006).

The development of polymerase chain reaction (PCR) sequence amplification coupled with increasingly inexpensive DNA sequencing technologies soon rendered RFLP mapping obsolete, and led to the development of remarkably efficient genotyping tools. Nonetheless, the RFLP maps that were generated in maize, tomato, wheat, barley and many other species permitted the chromosomal mapping of genes controlling economically important traits and led to the development of a genetic marker-assisted selection as a productive and practical plant breeding tool. Characterization of quantitative trait loci, genes impacting traits of interest in a quantitative rather than a qualitative Mendelian fashion, became a common practice in the 1990s. Several research groups were able to utilize genetic markers as starting points to identify and clone the genes impacting traits of interest, ranging from pathogen resistance to seed composition.

All plants derive from common ancestors, and structural similarities among genomes of relatively closely-related plant species are obvious. Among the *Triticeae*, seven pairs of mostly-metacentric chromosomes form the standard diploid genome. Barley and rye are diploids, with seven pairs of chromosomes. Durum wheat is a tetraploid, with fourteen

pairs of chromosomes and bread wheat is a hexaploid, with twenty one chromosome pairs. Synteny, the conserved physical relationship among genes on segments of chromosomes across related species, is generally observed across these species. When extended among to more distantly related rice and wheat or maize and rice, synteny can still be observed, although often gene locations have been altered by transposition or chromosome reorganization (Bennetzen *et al.*, 2005).

As DNA sequencing became less expensive and more efficient, sequencing genomes from a few plant species became an attractive and attainable goal. *Arabidopsis thaliana* was first sequenced; rice, grape and the model cool season grass *Brachypodium distachyon* soon followed. As the human genome program has demonstrated, within-species diversity is greater than we thought. Large-scale sequencing of the genomes of multiple individuals of important plant species will continue and will expand as sequencing costs decline. Although the cost of obtaining DNA sequence data continues to dramatically decline, assembling the relatively short DNA sequences in their proper order (this is called contig assembly) remains a daunting task. Since the genome of most crop plants contains transposons and retro-transposons that are imperfectly repeated thousands of times, it is difficult to know, when one sequences a retro-transposon, where in the genome it specifically belongs.

Several attempts have been made to estimate the number of genes expressed in the life of a plant. Estimates range from around 100,000 genes expressed during the lifespan of a tobacco plant to around 25,000 genes expressed in *Arabidopsis*. As the understanding of gene structure and regulation has advanced the interpretation of the plant transcriptome has evolved. Many research programs focused their effort on isolating messenger RNA from genes that were transcribed. These isolated mRNA molecules were copied to their complementary DNA molecules using the enzyme reverse transcriptase, and were then converted to double stranded DNA molecules which were then cloned and sequenced.

The relatively short (generally less than 1000 base pair) DNA sequences were called “expressed sequence tags” (ESTs) since the cloning process generally resulted in a truncated equivalent of the original mRNA molecule. EST cDNA sequencing projects have developed databases of sequences derived from isolated mRNA from many crops, with many plants providing hundreds of thousands of sequence files that have been condensed into contigs of related sequences. The barley (*Hordeum vulgare*) EST library contains more than 440,000 EST sequences that resolve into 28,000 contigs (<http://harvest.ucr.edu/>), each representing a gene or closely related family of genes.

Technologies permitting analysis of gene expression have advanced on pace with DNA sequence analysis techniques. While Northern blot analysis provided semi-quantitative analysis of gene expression, quantitative rt-PCR, microarrays, Serial Analysis of Gene Expression (SAGE), Illumina and Affymetrix technologies (see below) simultaneously permit analysis of most or all of the genes expressed in a tissue.

Among the most surprising discoveries in biology was the elucidation of small RNA mediated gene suppression, first discovered in plants (Herr and Baulcombe, 2004). In recent years, the RNA interference pathway has led to the development of novel

approaches to human disease management in addition to providing an entirely novel route by which we determine gene function.

As climate changes, the global population grows and competition for water resources increases, agricultural systems must become more productive and more efficient. Plant genetic resources provide still untapped reservoirs of genes that can contribute to better local adaptation and productivity. Plant genomics research provides insight into the genetic diversity that will be needed to prosper in the 21st century.

3. Structure and Composition of Plant Genomes

Plant genomes vary in genome size by a factor of 1000, but why they vary so dramatically is less obvious. Genome size variation appears to be poorly correlated with transcriptome complexity, and seems more dependent on the activity of transposable elements in the recent to distant past. Initially, researchers studying genome structure and composition relied on DNA re-naturation analysis. Typically, genomic DNA would be sheared to desired lengths by passage through a narrow orifice, appropriate size fractions would be denatured by heating and allowed to re-anneal under carefully controlled conditions. High copy number fragments would re-anneal more rapidly than would low copy number fragments, and single copy number fragments would find their complements and re-anneal last.

The rate of DNA re-annealing was monitored, and the relative proportions of DNA sequences of differing copy number determined. These analyses were represented as a C_0t curve. As with most of the first generation biotechnologies, the reliability of results depended on the skill and insight of the scientists performing the assays.

4. Mapping Plant Genomes

4.1. Mapping Technologies

It was well-understood that genes could be organized into chromosome-length linkage groups through genetic linkage analysis if scorable allelic variants for those genes could be found. Prior to 1980, those alleles typically altered the plant phenotype in a way that could be scored visually. Genes with these 'naked eye polymorphism' alleles were localized to positions on chromosomes through the use of cytogenetic lines like trisomics or chromosome addition or substitution lines.

Prior to 1980 each of the chromosomes of barley, maize, wheat and *Arabidopsis thaliana* contained some genetic markers, but the amount of work required to map each gene and the scarcity of new 'naked eye polymorphisms' precluded the development of densely-populated linkage maps. Once restriction fragment length polymorphisms (RFLPs) were suggested as a route to develop dense linkage maps in humans, the route to reasonably dense, informative linkage maps in plants became clear. RFLP maps in maize, barley, rice, tomato, pepper and potato were produced in the 1980s.

RFLPs provide robust and reliable data that permit unambiguous interpretation of allelic state information for an unlimited number of genetic polymorphisms. Unfortunately,

this technology is dependent on highly skilled technologists, the use of large quantities of genomic DNA, and consumption of enormous quantities of nucleotide tri-phosphates labeled with radioactive phosphorous. Mullis' discovery of the DNA polymerase chain reaction (PCR) changed genotype analysis virtually overnight. With RFLP analysis, the electrophoretic mobility was determined of a fragment of DNA that comprises 1/1,000,000th of the DNA in the electrophoresis sample. This requires the exquisite sensitivity provided by the use of radioactively labeled DNA probes. PCR allows to amplify specific DNA fragments millions-fold, and directly determines their structural characteristics. PCR is less expensive than RFLP analysis, does not demand the use of radioactive isotopes, and demands far less technical skill than does RFLP analysis.

In the late 1980s and early 1990s, many research groups determined that a subset of genes contained repeat sequences composed of copies of dimeric (e.g. AC) or trimeric (e.g. CAG) repeats (Akkaya *et al.*, 1992). During DNA replication, these simple repeat sequences tend to change in copy number, and if one uses PCR primers that surround one of these variable repeat sequences, allelic variation can be observed as fragment length polymorphisms by gel electrophoresis. Genes that contain these simple sequence repeats (SSRs) have been found in all surveyed eukaryotes, and these have proven to be extraordinarily useful genetic markers.

DNA sequencing costs declined as sequencing technology improved. While it was impractical to consider sequencing most crop genomes in the 1990's, sequencing portions of large numbers of genes was straightforward. Messenger RNA (mRNA) is easy to isolate from plant tissues, and converting mRNA to DNA is easily done using reverse transcriptase and DNA polymerase. Once cloned, these DNA fragments are easily sequenced, and represent the sequences of the expressed genes from a tissue.

The Institute for Genome Research (TIGR) was the initial repository for these Expressed Sequence Tag (EST) sequences, and by 2003 more than 3 million plant EST sequences had been reported. These have been utilized to enable the development of gene expression tools and massively parallel single nucleotide polymorphism (SNP) genotyping technologies (<http://www.illumina.com>). Table 1 lists some of the world's plant genome databases.

Name	Curators	Content
EMBL Nucleotide Sequence Database	http://www.ebi.ac.uk/embl/ European Bioinformatics Institute Hinxton, Cambridge, UK	Data and tools to analyze genes, gene expression, proteins, small molecules, pathways and much more.
The National Center for Biotechnology Information (NCBI)	www.ncbi.nlm.nih.gov National Library of Medicine, National Institutes of Health Beltsville, MD, USA	Genomes, nucleotide sequence information, pubmed, sequence analysis tools and more.
UK CropNet	ukcrop.net IGER, JIC, NASC and SCRI Six projects in four institutes in the United Kingdom	<i>Arabidopsis</i> , barley, Brassica, forage grasses, and millet databases and comparative analysis tools.
National BioResource Project (NBRP)	http://www.nbrp.jp A collaboration of many Japanese scientific institutes.	Resource center and repository for genomics projects.

The IPK Crop EST Database (CR-EST)	ipk-gatersleben.de Leibniz Institute of Plant Genetics and Crop Plant Research Gatersleben, Germany	EST databases for wheat, potato, barley, pea, petunia and tobacco.
AFRICANCROPS.NET	www.africancrops.net Alliance for a Green Revolution in Africa. Rockefeller Foundation, Bill and Melina Gates Foundation	African crop breeding networks, databases, seed source, news, free access portals, training programs.
J. Craig Venter Institute	www.jcvi.org J. Craig Venter Institute Rockville, MD, USA	<i>Brassicaceae</i> , <i>Medicago</i> , castor bean, rice genomics and much more.
Harvest	harvest.ucr.edu University of California Riverside, CA, USA	EST datasets and tools for barley, rice, citrus, cowpea, coffee, <i>Brachypodium</i> , wheat and soybean.
TAIR	www.arabidopsis.org The Arabidopsis Information Resource Carnegie Inst. Stanford, CA, USA	Complete genome browser, extensive suite of tools and accession to genetic stocks.
OryzaBase	www.shigen.nig.ac.jp/rice/oryzabase National Institute of Genetics in Japan	Integrated rice science, genetic map and mutant database
Gramene	http://www.gramene.org USDA-ARS, Cornell University Cold Spring Harbor, NY, USA	Comparative genomic analysis in the grasses with an emphasis on rice.
Graingenes	wheat.pw.usda.gov USDA, ARS, Albany, CA Cornell Univ., Ithaca, NY, USA	Genetic, genomic and phenotypic information for the Triticeae and <i>Avena</i> .
Plant GDB	plantgdb.org Iowa State University Ames, IA, USA	Maize EST database, tools and resources for plant genomics.
Panzea	panzea.org USDA-ARS Cornell University, Ithaca, NY, USA	Maize and teocinte genome data and software for genome and association analysis.
Soybase	soybase.org USDA-ARS Iowa State University, Ames, IA	Genetics data and tools for soybean and legumes. Breeders Toolbox.
CottonDB	cottondb.org USDA-ARS College Station, Texas, USA	Cotton Genome Browser, genes, genetics and taxonomy.

Table 1. Some of the Many Plant Genome Databases containing plant genomics data.

-
-
-

TO ACCESS ALL THE 23 PAGES OF THIS CHAPTER,
Visit: <http://www.eolss.net/Eolss-sampleAllChapter.aspx>

Bibliography

Akkaya, M.S., Bgwat, A.A. and Cregan, P.B. (1992). *Length Polymorphisms of Simple Sequence Repeat DNA in Soybean*. *Genetics*, 132: 1131-1139. [Perry Cregan's group pioneered the use of microsatellites in this paper].

Alwine, J., Kemp, D. and Stark, G.R. (1977). *Method for Detection of Specific RNAs in Agarose Gels by Transfer to Diazobenzyloxymethyl-paper and Hybridization with DNA Probes*. *Proc. Natl. Acad. Sci. USA*, 74: 5350-5354. [This paper provides a description of northern blots, the cornerstone of gene expression research].

Barrett, T., Troup, D.B., Wilhite, S.E., Ledoux, P., Rudnev, D., Evangelista, C., Kim, I.F., Soboleva, A., Tomashevsky, M. and Edgar, R. (2006). *NCBI GEO: Mining Tens of Millions of Expression Profiles — Database and Tools Update*. *Nucl. Acids. Res.*, 35: D760-D765. [This paper provides a comprehensive review of genome databases.]

Baulcombe, D.C. and Kay, J.L. (1980). *Polyadenylated RNA Sequences which are Reduced in Concentration following Auxin Treatment of Soybean Hypocotyls*. *J. Biol. Chem.*, 255: 8907-8913. [This was one of the first explorations of transcriptional gene control in plants.]

Bennetzen J.L., Ma J. and Devos, K.M. (2005). *Mechanisms of Recent Genome Size Variation in Flowering Plants*. *Annals of Botany*, 95(1): 127-132. [This is an excellent review of plant genome structure and variation.]

Botstein, D., White, R.L., Skolnick, M. and Davis, R.W. (1980). *Construction of a Genetic Linkage Map in Man Using Restriction Fragment Length Polymorphisms*. *Amer. J. Hum. Genet.*, 32: 314-331. [This paper created the field of eukaryotic restriction fragment length polymorphism mapping.]

Hayes, P.M., Liu, B.H., Knapp, S.J., Chen, F., Jones, B., Blake, T., Frankowiak, J., Rasmussen, D., Sorrells, M., Ullrich, S.E., Wesenberg, D. and Kleinjohs, A. (1994). *Quantitative Trait Locus Effects and Environmental Interaction in a Sample of North American Barley Germplasm*. *Theor. Appl. Genet.*, 87: 397-401. [In this paper, a broad coalition of researchers applied genome mapping techniques to map the locations of genes that contribute to plant performance.]

Heil, M. and Ton, J. (2008). *Long-distance Signaling in Plant Defense*. *Trends Plant Sci*, 13: 264-272. [This is an excellent review of plant defense responses.]

Herr, A.J. and Baulcombe, D.C. (2004). *RNA Silencing Pathways in Plants*. *Cold Spring Harb. Symp. Quant. Biol.*, 69: 363-370. [An excellent review of siRNA-mediated gene regulation in plants.]

Jones, D.F. (1917). *Dominance of Linked Factors as a Means of Accounting for Heterosis*. *Proc. Nat. Adac. Sci. (USA)*, 3: 310-312. [This is a wonderful, small paper that provided an explanation of heterosis.]

Kalantidis, K., Schumacher, H.T., Alexiadis, T. and Helm, J.M. (2008). *RNA Silencing Movement in Plants*. *Biology of the Cell*, 100: 13-26. [This is a recent review of RNA silencing and signal transduction.]

Lashkari, D.A., DeRisi, J.L., McCusker, J.H., Namath, A.F., Gentile, C., Hwang, S.Y., Brown, P.O. and Davis, R.W. (1997). *Yeast Microarrays for Genome Wide Parallel Genetic and Gene Expression Analysis*. *Proc. Natl. Acad. Sci. USA*, 94: 13057–13062. [This was the seminal paper demonstrating the value of microarrays in monitoring gene expression.]

McClintock, B. (1987). *The Discovery and Characterization of Transposable Elements: The Collected Papers of Barbara McClintock*. Garland Publications, New York. [Barbara McClintock was the most insightful geneticist of her generation. These are her collected works.]

Sears, E.R. (1969). *Wheat Cytogenetics*. *Ann. Rev. Genet.*, 3: 451-468. [This paper shows how Ernie Sears and his colleagues developed the cytogenetic tools that permitted initial gene mapping in wheat.]

Shena, M., Shalon, D., Davis, R.W., Brown, P.O. (1995). *Quantitative Monitoring of Gene Expression Patterns with a Complementary DNA Microarray*. *Science*, 270: 467-470. [This short paper provides a graphic illustration of the use microarray technology to evaluate gene expression.]

Tang, G (2005). *siRNA and miRNA: An Insight into RISCs*. Trends Biochem Sci., 30: 106-114. [This is an excellent review explaining mechanisms underlying small RNA mediated gene suppression.]

Tanksley, S.D. and Fulton, T.M. (2006). *Dissecting Quantitative Trait Variation: Examples from the Tomato*. Euphytica, 154(3): 365-370. [This is an excellent, insightful review of quantitative trait locus analysis.]

The *Arabidopsis* genome initiative, (2000). *Analysis of the Genome Sequence of the Flowering plant Arabidopsis thaliana*. Nature, 408: 796-815. [This paper reports the first whole genome sequence of a plant.]

Velculescu, V. E., Zhang, L., Vogelstein, B. and Kinzler, K. W. (1995). *Serial Analysis Of Gene Expression*. Science, 270: 484-487. [This paper explains how SAGE works.]

Zipper, H., Brunner, H., Bernhagen, J. and Vitzthum, F. (2004). *Investigations on DNA Intercalation and Surface Binding by SYBR Green I: Its Structure Determination and Methodological Implications*. Nucl. Acids Res., 32: 103. [This paper provided the basis for quantitative PCR analysis, one of the primary technologies used to evaluate transcriptional gene regulation.]

Biographical Sketches

Thomas Blake was raised on an olive orchard in Northern California. He received his B.S. in Genetics from the University of California (Davis) in 1976, his M.S. (agronomy) from South Dakota State University in 1979, and his Ph. D. from Washington State University (Genetics) studying under Dr. R.A. Nilan in 1983. He began mapping genes as a graduate student, and focused on mapping genes contributing to traits of agricultural importance when he joined the faculty of Montana State University as barley breeder and geneticist in 1984.

Dr. Blake took a 2-year leave from Montana State University to serve as Director of the Germplasm Program at the International Center for Agricultural Research in the Dry Areas (ICARDA) in 2002 and 2003. He returned to Montana State University in January, 2004, and is currently releasing drought tolerant 2-rowed malt and feed barley varieties.

Victoria Carollo Blake grew up in the San Francisco Bay Area. She received her B.S. in Biology from San Jose State University and her Ph.D. in Plant Biology from U.C. Davis. During her graduate studies in the Viticulture and Enology Department to characterize malic enzyme gene expression in ripening fruit, she would often spend fall semesters working the 'crush' in the Napa Valley for practical agricultural experience. She then took a Molecular Biologist position at the USDA ARS Western Regional Research Center in Albany, CA, to work as a curator for GrainGenes, the international database for the Triticeae and Avena.

In 2006, Victoria moved to Montana to join Tom Blake's lab and family in Bozeman. Her part of the barley project is to identify barley varieties whose forage and straw would provide a good feedstock for biofuel. They are making great progress in this area and hope to have a novel, inexpensive and environmentally friendly form of ethanol production from barley straw available by 2011.

Jackie Campbell received her B.S. degree (Biology) from the University of Colorado in 2004, and her M.S. degree from Montana State University in 2007, studying with Dr. Mike Giroux. Ms. Campbell has won several awards, including the Presidential Scholarship for Academic Excellence (University of Colorado), the Mildred Livingston Presidential Scholarship (Montana State University) and the Bayard Taylor Award for Outstanding Graduate Students (Montana State University).

Ms. Campbell is currently pursuing her Ph. D. (plant genetics) with Dr. Li Huang in the Plant Sciences and Plant Pathology Department, Montana State University. Her research topic is gene regulation, utilizing Barley Stripe Mosaic Virus as a vector for siRNA-mediated gene regulation.

Plant Transcriptomes

A plant transcriptome refers to the complete set of RNA molecules transcribed from the genes of a plant species at a specific time or under specific conditions. It represents the dynamic expression profile of genes in a plant and provides valuable insights into gene function, regulation, and the plant's response to various environmental stimuli.

Plant transcriptomics involves the large-scale analysis of the transcriptome using techniques such as microarrays and RNA sequencing (RNA-seq). By studying the transcriptome, researchers can identify and quantify the RNA molecules present in a plant, including protein-coding transcripts, non-coding RNAs, and splice variants. This information helps in understanding the genetic basis of plant traits and provides a comprehensive view of gene expression patterns.

The transcriptome analysis provides several benefits and applications in plant biology:

Gene Expression Profiling: Transcriptomics enables the study of gene expression patterns across different tissues, developmental stages, and environmental conditions. It allows the identification of genes that are activated or repressed in response to specific stimuli, facilitating the understanding of plant development, physiological processes, and stress responses.

Discovery of Novel Genes and Splice Variants: Transcriptome analysis aids in the identification of novel genes that may not be annotated in the plant genome. It also helps in detecting alternative splicing events, where different combinations of exons are joined together to produce multiple transcript isoforms. This knowledge enhances our understanding of gene structure and function.

Functional Annotation and Gene Regulatory Networks: By correlating transcriptomic data with functional information, researchers can assign putative functions to genes of interest. Transcriptomics also enables the reconstruction of gene regulatory networks by identifying co-

expressed genes and transcription factor-target relationships. These networks provide insights into the complex regulatory mechanisms controlling plant development and responses to environmental cues.

Comparative Transcriptomics: Comparative transcriptomics involves the comparison of transcriptomes across different plant species or genotypes. It helps identify conserved and species-specific gene expression patterns, facilitating the understanding of evolutionary relationships and functional conservation/divergence among plant species.

Marker Development and Breeding: Transcriptomic data can be utilized to develop molecular markers, such as single nucleotide polymorphisms (SNPs), that are associated with traits of interest. These markers aid in marker-assisted selection (MAS) and genomic selection (GS), facilitating crop improvement programs and accelerating breeding processes.

Genetic Engineering and Biotechnology: Transcriptome analysis is essential for understanding the expression patterns of genes involved in metabolic pathways and desired traits. This information guides the engineering of plants through genetic modification or genome editing techniques, enabling the production of plants with improved agronomic traits, enhanced nutritional content, or resistance to pests and diseases.

In summary, plant transcriptomics plays a crucial role in unraveling the complexity of gene expression in plants. It provides insights into gene function, regulatory mechanisms, and plant responses to environmental stimuli. The knowledge gained from transcriptome analysis has broad applications in agriculture, crop improvement, and biotechnology, contributing to advancements in plant science and the development of sustainable agricultural practices.

Plant small RNAs

Plant small RNAs (sRNAs) are short RNA molecules that play crucial roles in regulating gene expression and controlling various biological processes in plants. These sRNAs are typically 20-24 nucleotides in length and are generated from longer RNA precursors through a process called RNA silencing or RNA interference (RNAi).

There are several types of plant sRNAs, each with its specific functions and biogenesis pathways:

MicroRNAs (miRNAs): MiRNAs are endogenous sRNAs that regulate gene expression by targeting messenger RNAs (mRNAs) for degradation or translational repression. They are generated from single-stranded RNA precursors called primary miRNAs (pri-miRNAs) and undergo a series of processing steps to form mature miRNAs. MiRNAs are involved in diverse processes such as development, growth, and response to environmental stresses.

Small interfering RNAs (siRNAs): SiRNAs are another class of endogenous sRNAs that play a vital role in defense against viral infections and transposable elements. They are derived from long double-stranded RNA molecules and can be further classified into two main types: trans-acting siRNAs (tasiRNAs) and heterochromatic siRNAs (hcsiRNAs). TasiRNAs are involved in the regulation of developmental processes, while hcsiRNAs are associated with heterochromatin formation and gene silencing.

Natural antisense transcript-derived siRNAs (nat-siRNAs): Nat-siRNAs are sRNAs derived from the complementary regions of natural antisense transcripts (NATs). NATs are RNA molecules transcribed from the opposite strand of a gene and can form double-stranded RNA structures, which are then processed into nat-siRNAs. These sRNAs have been implicated in gene regulation, stress responses, and genome stability.

Phased secondary siRNAs (phasiRNAs): PhasiRNAs are a specific class of siRNAs that are generated in a phased manner from noncoding RNA transcripts targeted by miRNAs. They are involved in plant reproductive development, including pollen and ovule development.

Long noncoding RNA-derived siRNAs (lsiRNAs): LsiRNAs are sRNAs derived from long noncoding RNA (lncRNA) precursors. These sRNAs are generated through the activity of RNA-dependent RNA polymerases (RDRs) and are involved in transcriptional gene silencing and chromatin modifications.

Plant sRNAs act as important regulators of gene expression by directing sequence-specific silencing of target genes. They mediate a wide range of biological processes, including development, stress responses, immune defense, and genome stability. Understanding the diverse functions and regulatory mechanisms of plant sRNAs provides valuable insights into the complexity of plant biology and offers potential applications in crop improvement and biotechnology.

Plant epigenomes

Plant epigenomes refer to the heritable modifications in gene expression that do not involve changes in the DNA sequence itself. These modifications occur at the epigenetic level and can influence gene activity and phenotype. Epigenetic changes play a vital role in regulating various aspects of plant development, stress responses, and adaptation to the environment.

The main epigenetic modifications in plants include DNA methylation, histone modifications, and small RNA-mediated gene silencing. Here is a brief overview of each:

DNA methylation: DNA methylation involves the addition of a methyl group to the cytosine residues of DNA molecules. In plants, DNA methylation occurs in three contexts: CG, CHG, and CHH (where H represents any nucleotide other than G). CG methylation is relatively stable and prevalent in the plant genome. CHG and CHH methylation, on the other hand, are established and maintained through the activity of different DNA methyltransferases. DNA methylation can regulate gene expression by blocking the binding of transcription factors or recruiting proteins that modify chromatin structure.

Histone modifications: Histones are proteins that form the backbone of chromatin, and their post-translational modifications can influence gene expression. Histone modifications include acetylation, methylation, phosphorylation, ubiquitination, and more. For instance, histone acetylation is generally associated with active gene expression, while methylation can be associated with either gene activation or repression depending on the context and specific residues modified. Histone modifications can alter chromatin structure and accessibility, thereby affecting gene expression patterns.

Small RNA-mediated gene silencing: Small RNAs, such as microRNAs and small interfering RNAs, play a crucial role in gene regulation through RNA silencing or RNA interference mechanisms. Small RNAs can guide sequence-specific degradation or translational repression of

target messenger RNAs (mRNAs), leading to gene silencing. These small RNAs are generated through various pathways and often act in a feedback loop to regulate their own biogenesis.

Plant epigenomes can be influenced by internal factors, such as developmental cues, and external factors, including environmental stresses and nutrient availability. Epigenetic modifications are mitotically and meiotically heritable, meaning they can be passed on to subsequent generations. This heritability allows plants to adapt and respond to changing environmental conditions and transmit epigenetic information to their offspring.

Studying plant epigenomes has become an important area of research in understanding plant biology and improving crop traits. By deciphering the epigenetic modifications and mechanisms involved, scientists can potentially develop strategies to manipulate plant epigenomes for crop improvement, stress tolerance, and disease resistance. However, further research is still needed to fully unravel the complexity of plant epigenetic regulation and its implications in plant biology and agriculture.